

AG
T

*Algebraic & Geometric
Topology*

Volume 21 (2021)

2–Segal objects and the Waldhausen construction

JULIA E BERGNER
ANGÉLICA M OSORNO
VIKTORIYA OZORNOVA
MARTINA ROVELLI
CLAUDIA I SCHEIMBAUER



2–Segal objects and the Waldhausen construction

JULIA E BERGNER
 ANGÉLICA M OSORNO
 VIKTORIYA OZORNOVA
 MARTINA ROVELLI
 CLAUDIA I SCHEIMBAUER

In a previous paper, we showed that a discrete version of the S_\bullet –construction gives an equivalence of categories between 2–Segal sets and augmented stable double categories. Here, we generalize this result to the homotopical setting, by showing that there is a Quillen equivalence between a model category for 2–Segal objects and a model category for augmented stable double Segal objects which is given by an S_\bullet –construction. We show that this equivalence fits together with the result in the discrete case and briefly discuss how it encompasses other known S_\bullet –constructions.

18D05, 18G55, 19D10, 55U35, 55U40; 18G30, 55U10

Introduction	1267
1. Summary of background results	1271
2. The generalized S_\bullet –construction	1285
3. Relationship with other S_\bullet –constructions	1297
4. The model structures	1305
5. The generalized S_\bullet –construction as a right Quillen functor	1309
6. The Quillen equivalence	1318
7. Variants of the model structures	1322
References	1324

Introduction

Waldhausen’s S_\bullet –construction, first described in [32], provides a way to define algebraic K –theory via classifying spaces of categories of certain diagrams, so that its output

is a simplicial space. While this construction was defined in the more general setting of categories with cofibrations and weak equivalences (often now called Waldhausen categories) it provides a different way to think about the algebraic K -theory of exact categories, originally constructed by Quillen [29].

Independently, Dyckerhoff and Kapranov [11] and Gálvez-Carrillo, Kock and Tonks [16] showed that the simplicial spaces obtained as the output of an S_\bullet -construction for exact categories have the additional structure of a 2-Segal space or *decomposition space*. It is particularly interesting that both sets of authors identify the output of the Waldhausen construction as a crucial example, although their approaches to this structure are quite different. The main starting point for our work was the following question:

Question *Is this source of examples exhaustive? In other words, does every 2-Segal space arise from such a construction?*

The main goal of this paper is to give a positive answer to this question for a generalization of the S_\bullet -construction.

To give a brief overview, first recall that a Segal space is a simplicial space X such that the Segal maps

$$X_n \rightarrow \underbrace{X_1 \times_{X_0}^h \cdots \times_{X_0}^h X_1}_n$$

are weak equivalences for $n \geq 2$. We can think of a Segal space X as having a space of objects X_0 , a space of morphisms X_1 , and up-to-homotopy composition which can be defined by the span

$$X_1 \times_{X_0}^h X_1 \xleftarrow{\cong} X_2 \rightarrow X_1,$$

since the first arrow is a weak equivalence. In other words, a Segal space is a topological category up to homotopy.

As a generalization, a 2-Segal space is a simplicial space X such that certain maps

$$X_n \rightarrow \underbrace{X_2 \times_{X_1}^h \cdots \times_{X_1}^h X_2}_{n-1}$$

are weak equivalences for $n \geq 3$. In this setting, we still have a space of objects X_0 and a space of morphisms X_1 , but we need not have composition of all possibly composable pairs of morphisms, since the first map in the span

$$X_1 \times_{X_0}^h X_1 \leftarrow X_2 \rightarrow X_1$$

is no longer necessarily invertible, even up to homotopy. However, we can think of a 2–Segal space as having a multivalued composition, where an element of $X_1 \times_{X_0} X_1$ can be lifted to any preimage in X_2 , which is in turn sent to its image in X_1 . Thus, two potentially composable morphisms could have no composite at all (if the preimage in X_2 is empty) or multiple composites (if the preimage has multiple elements). The homotopy invertibility of the 2–Segal maps given above is used to prove that this multivalued composition is homotopy associative. We thus think of the structure of a 2–Segal space as that of a category with multivalued composition up to homotopy.

In [11], Dyckerhoff and Kapranov include an additional assumption, and call a 2–Segal space *unital* if composition with identity morphisms always exists and is unique up to homotopy. However, a recent surprising result of Feller, Garner, Kock, Underhill-Proulx and Weber [12] shows that every 2–Segal space is unital. We thus frequently omit the term “unital” unless we find it useful for clarification.

In previous work [6] we considered the motivating question above in a discrete setting, in which the output was a simplicial set, rather than a simplicial space. We gave a positive answer, in that we showed that the category of 2–Segal sets is equivalent to the category of *augmented stable double categories*; the equivalence is given by a discrete S_\bullet –construction and its inverse by a path construction. While illuminating in its own right, this discrete setting is insufficient to encompass even the classical example of exact categories, for which we need to work homotopically.

In particular, we expect not an equivalence of categories, but instead an equivalence of homotopy theories, given by a Quillen equivalence of model categories. Indeed, we need not restrict ourselves to working solely in the context of spaces, but can consider 2–Segal objects in any sufficiently nice model category \mathcal{C} . By a result of Dyckerhoff and Kapranov, there is a model structure on the category of simplicial objects in \mathcal{C} in which the fibrant objects are the 2–Segal objects.

To make our comparison, then, we first need a model category whose fibrant objects are augmented stable double Segal objects in \mathcal{C} , the homotopical analogues of augmented stable double categories. While double Segal objects give a homotopical generalization of double categories, we can also describe the appropriate generalizations of the notions of augmentation and stability in that context. The resulting structures can be realized as the fibrant objects in a model structure on a category of bisimplicial objects in \mathcal{C} with additional structure.

We now state the main result of this paper, which is stated more precisely in [Theorem 6.1](#):

Theorem *There is a Quillen equivalence between the model category for 2–Segal objects and the model category for augmented stable double Segal objects.*

As in the discrete case, the functor from augmented stable double Segal objects can be regarded as a version of the S_\bullet –construction, and its adjoint can be thought of as a path construction. We show in [Section 3.1](#) that the restriction to the discrete case recovers the functors from our previous work.

A topic of much recent work has been the generalization of the classical S_\bullet –construction to more general contexts, such as for stable $(\infty, 1)$ –categories and (proto-)exact $(\infty, 1)$ –categories, for example by Barwick [\[5\]](#), Blumberg, Gepner and Tabuada [\[8\]](#) and Fiore and Pieper [\[15\]](#). In a companion paper [\[7\]](#) we show that our construction recovers and generalizes these results; we give a summary in [Section 3.2](#).

Outline of the paper

In [Section 1](#), we recall some background information about enriched localizations and 2–Segal spaces, as well as summarize our previous results in the discrete setting. [Section 2](#) is concerned with developing the definition of augmented stable double Segal objects and defining the S_\bullet –construction in this context. In [Section 3](#) we discuss how this construction is a generalization of previously known ones: in [Section 3.1](#) we show explicitly that we recover the one from the discrete setting, and in [Section 3.2](#) we summarize how it generalizes the S_\bullet –construction as defined for proto-exact $(\infty, 1)$ –categories.

We turn to developing the model structures that we need in [Section 4](#). We show that there is a Quillen pair between these model structures in [Section 5](#), and in [Section 6](#) we show that it is a Quillen equivalence.

Finally, in [Section 7](#) we consider some variants of the model structures and comparisons that we have developed in this paper.

Acknowledgements

We would like to thank the organizers of the Women in Topology II Workshop and the Banff International Research Station for providing a wonderful opportunity for collaborative research. Conversations with T Dyckerhoff, I Gálvez–Carrillo, J Kock, L Meier and A Tonks were helpful. We also thank the referee for helpful comments and suggestions.

Bergner was partially supported by NSF CAREER award DMS-1659931. Osorno was partially supported by a grant from the Simons Foundation (#359449, Angélica Osorno), the Woodrow Wilson Career Enhancement Fellowship and NSF grant DMS-1709302. Rovelli and Scheimbauer were partially funded by the Swiss National Science Foundation, grants P2ELP2_172086 and P300P2_164652. Scheimbauer was partially funded by the Trond Mohn Foundation and would like to thank the University of Oxford for its hospitality. The authors would like to thank the Isaac Newton Institute for Mathematical Sciences, Cambridge, for support and hospitality during the program *Homotopy harnessing higher structures*, where work on this paper was undertaken. This work was supported by EPSRC grant no. EP/K032208/1.

1 Summary of background results

In this section we recall some necessary background material. First, we review some standard results concerning presheaves in a symmetric monoidal model category. We then review the theory of 2–Segal objects, which are central to our work here, and summarize the comparison between 2–Segal sets and augmented stable double categories.

1.1 Enriched model structures on presheaf categories

We begin with a review of the basic theory of presheaf categories in a symmetric monoidal category and some relevant results in the context of model categories. One purpose of this section is to set notation that we use throughout; the reader may choose to skip this section and refer back as needed.

Let I be a small category and $\mathcal{C} = (\mathcal{C}, \otimes, \mathbb{1})$ a complete and cocomplete closed symmetric monoidal category. We denote by $\mathrm{Fun}(I, \mathcal{C})$ the category of functors $I \rightarrow \mathcal{C}$. As explained in [25, Section 2.5], there is an adjunction

$$d : \mathcal{S}et \rightleftarrows \mathcal{C} : U$$

between the functors defined by

$$d(X) := \coprod_X \mathbb{1} \quad \text{and} \quad U(Y) := \mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, Y).$$

Objects in the image of d are called *discrete*. Here, and elsewhere in the paper, we follow the convention of displaying the left adjoint topmost.

The following proposition enables us to view every presheaf in $\mathcal{S}et$ as a discrete presheaf in \mathcal{C} via the functor d_* :

Proposition 1.1 *The adjunction $d: \mathcal{S}et \rightleftarrows \mathcal{C}: U$ induces an adjunction*

$$d_*: \mathbf{Fun}(I, \mathcal{S}et) \rightleftarrows \mathbf{Fun}(I, \mathcal{C}): U_*.$$

For simplicity, we frequently omit the notation d_* when referring to discrete presheaves.

An important feature of the category $\mathbf{Fun}(I, \mathcal{C})$ is that it has a canonical enrichment.

Proposition 1.2 [25, Section 2.2] *The category $\mathbf{Fun}(I, \mathcal{C})$ is enriched over \mathcal{C} . Given objects X and Z of $\mathbf{Fun}(I, \mathcal{C})$, the mapping object in \mathcal{C} is the end*

$$\mathrm{Map}_{\mathbf{Fun}(I, \mathcal{C})}(X, Z) := \int_I \mathrm{Map}_{\mathcal{C}}(X_i, Z_i).$$

Now suppose that $f: I \rightarrow J$ is a functor between small categories. The associated precomposition functor $f^*: \mathbf{Fun}(J, \mathcal{C}) \rightarrow \mathbf{Fun}(I, \mathcal{C})$ admits a right adjoint $f_*: \mathbf{Fun}(I, \mathcal{C}) \rightarrow \mathbf{Fun}(J, \mathcal{C})$ via right Kan extension. The following statement is a variant of [25, Theorem 4.50]:

Proposition 1.3 *The precomposition functor f^* and its right adjoint f_* form a \mathcal{C} -enriched adjunction*

$$f^*: \mathbf{Fun}(J, \mathcal{C}) \rightleftarrows \mathbf{Fun}(I, \mathcal{C}): f_*.$$

Now we equip \mathcal{C} with a model structure that is compatible with the closed symmetric monoidal structure.

Assumption 1.4 Let $\mathcal{C} = (\mathcal{C}, \otimes, \mathbb{1})$ be a combinatorial closed symmetric monoidal model category in which all objects are cofibrant, so in particular it is left proper.

Examples include the category $s\mathcal{S}et$ of simplicial sets endowed with the classical model structure due to Quillen [28], the category $\mathcal{G}pd$ of groupoids endowed with the canonical model structure [1, Section 5], and the category $\mathcal{S}et$ endowed with the (unique) model structure in which weak equivalences are isomorphisms.

Remark 1.5 It is possible that our results can be extended to more general model structures, but these assumptions are convenient for our arguments. In particular, the assumption that all objects are cofibrant appears centrally in two places. First, we use the condition that the unit object is cofibrant when comparing discrete objects to objects in \mathcal{C} , for example already in the next lemma. Second, we use that all objects are cofibrant in establishing the Quillen adjunction in Proposition 5.2, since the left adjoint there need not be derived.

Ideally we would like to have similar results in more generality. For example, the usual model structure on topological spaces does not have all objects cofibrant. Another example is the model category of derived stacks from [31], which appeared in the context of 2–Segal spaces in [11]; but even affine stacks need not be cofibrant. However, we expect our results could be adjusted to this setting.

The following lemma incorporates the appropriate model structures to give us more information about the adjunction from Proposition 1.1. We leave the proof to the reader.

Lemma 1.6 *The functor*

$$d_* : \text{Fun}(I, \text{Set}) \rightarrow \text{Fun}(I, \mathcal{C})$$

sends monomorphisms to levelwise cofibrations.

Motivated by this lemma, we consider the category of presheaves on \mathcal{C} with the injective model structure, in which both the weak equivalences and the cofibrations are given levelwise.

Theorem 1.7 [3, Propositions 4.50 and 4.51] *The category $\text{Fun}(I, \mathcal{C})$ admits the injective model structure. This model structure is combinatorial, symmetric monoidal, enriched over \mathcal{C} with respect to the mapping spaces from Proposition 1.2, and has all objects cofibrant.*

Throughout this paper, unless indicated otherwise, we assume that $\text{Fun}(I, \mathcal{C})$ is equipped with the injective model structure. When we want to emphasize this structure or when there might be ambiguity, we denote this model structure by $\text{Fun}(I, \mathcal{C})_{\text{inj}}$. We have chosen to use the injective model structure because all objects are cofibrant, and hence derived mapping objects can be computed easily, as the following proposition demonstrates. We discuss other model structures on our categories of interest in Section 7.

Definition 1.8 Given objects X and Z of $\text{Fun}(I, \mathcal{C})$, the *derived mapping object* is given by

$$\text{Map}^h(X, Z) := \text{Map}(X, Z^f),$$

where $(-)^f$ denotes a functorial fibrant replacement in the injective model structure on $\text{Fun}(I, \mathcal{C})$.

More information about derived mapping objects and enriched model categories can be found in [3]. We collect the main properties in the following proposition:

Proposition 1.9 *The derived mapping object is functorial, respects weak equivalences in both variables and coincides up to equivalence with the strict mapping space when the second variable is fibrant.*

We now recall some terminology and the construction of enriched localizations of a model category.

Definition 1.10 Let S be a set of maps in $\text{Fun}(I, \mathcal{C})$ with the injective model structure.

- An object X of $\text{Fun}(I, \mathcal{C})$ is S -local if it is fibrant and, for every map $f: A \rightarrow B$ in S , the induced map

$$f^*: \text{Map}^h(B, X) \rightarrow \text{Map}^h(A, X)$$

is a weak equivalence in \mathcal{C} .

- A map $g: C \rightarrow D$ in $\text{Fun}(I, \mathcal{C})$ is an S -local equivalence if for every S -local object X , the induced map

$$g^*: \text{Map}^h(D, X) \rightarrow \text{Map}^h(C, X)$$

is a weak equivalence in \mathcal{C} .

Since we use the injective model structure in which all objects are cofibrant, and S -local objects are assumed to be fibrant, the underived mapping spaces in our setting model derived mapping spaces, and hence can be used in this definition.

With these definitions in hand, we can define the enriched localization of the injective model structure on $\text{Fun}(I, \mathcal{C})$ with respect to the maps in S .

Theorem 1.11 [3, Theorem 4.46] *Let S be a set of maps of $\text{Fun}(I, \mathcal{C})$. There exists a cofibrantly generated model structure on $\text{Fun}(I, \mathcal{C})$, denoted by $\text{Fun}(I, \mathcal{C})_S$, in which*

- *the cofibrations are the levelwise cofibrations, and in particular all objects are cofibrant;*
- *the fibrant objects are the S -local objects;*
- *the weak equivalences are the S -local equivalences; and*
- *the weak equivalences between fibrant objects are the levelwise weak equivalences.*

The definitions we have used, and the previous theorem, hold for more general model categories than the injective model structure, in particular for the projective model structure on $\text{Fun}(I, \mathcal{C})$, which we consider in [Section 7](#). The S –local objects are dependent on the model structure chosen, but the S –local equivalences only depend on the weak equivalences thereof.

1.2 2–Segal objects

In the context of a category \mathcal{C} satisfying [Assumption 1.4](#), we recall the notion of a Segal map.

Notation 1.12 Let $n \geq 0$. Denote by $\mathcal{I}[n]$ the spine of the standard n –simplex, ie the simplicial set which is the colimit of standard 1–simplices

$$\Delta[1] \amalg_{\Delta[0]} \cdots \amalg_{\Delta[0]} \Delta[1] \cong \mathcal{I}[n] \subset \Delta[n],$$

where the image of the i^{th} copy of $\Delta[1]$ is the simplex of $\Delta[n]$ with vertices i and $i + 1$.

Definition 1.13 A *Segal object* in \mathcal{C} is a simplicial object in \mathcal{C} such that, for every $n \geq 2$, the Segal map

$$X_n \rightarrow \underbrace{X_1 \times_{X_0} \cdots \times_{X_0} X_1}_n \simeq \text{Map}_{\mathcal{C}^{\Delta^{\text{op}}}}^h(\mathcal{I}[n], X)$$


which is induced by the inclusion $\mathcal{I}[n] \hookrightarrow \Delta[n]$ is a weak equivalence.

Dyckerhoff and Kapranov define 2–Segal objects using higher-dimensional analogues of these Segal maps, in the sense that they arise from polygonal decompositions of polygons [\[11\]](#).

For an $(n+1)$ –gon in the plane with vertices labeled cyclically by $\{0, 1, \dots, n\}$, we consider its polygonal decompositions with vertices chosen amongst the vertices of the polygon. Thus, a polygonal decomposition consists of a collection of noncrossing diagonals subdividing the $(n+1)$ –gon into other polygons. For technical reasons, we view such a decomposition as the collection of the resulting polygons together with the defining diagonals.

Formally, we can regard such a decomposition as a subposet P of the power set of $\{0, 1, \dots, n\}$ which is closed under nonempty intersection. For instance, the polygonal

decomposition of the square



corresponds to the poset $\{\{1, 3\}, \{1, 2, 3\}, \{0, 1, 3\}\}$.

Definition 1.14 Let P be a polygonal decomposition of a regular $(n+1)$ –gon, consisting of $(k+1)$ –gons of the form $\{i_0, \dots, i_k\}$ for varying k . The P –Segal map of a simplicial object X in \mathcal{C} is the induced map

$$X_n \rightarrow \operatorname{holim}_{\{i_0, \dots, i_k\} \in P} X_k,$$

where the homotopy limit is taken over the poset of polygons $\{i_0, \dots, i_k\}$ occurring in the decomposition P , ordered by inclusion of vertices.

Now we are able to define 2–Segal objects, following [11]. Here we use the original definition, given in terms of triangulations, or polygonal decompositions into triangles.

Definition 1.15 A 2–Segal object in \mathcal{C} is a simplicial object in \mathcal{C} such that, for every $n \geq 3$ and every triangulation T of a regular $(n+1)$ –gon, the T –Segal map

$$X_n \rightarrow \operatorname{holim}_{\substack{\{i_0, \dots, i_k\} \in T \\ k=1,2}} X_2 \simeq \underbrace{X_2 \overset{h}{\times}_{X_1} \cdots \overset{h}{\times}_{X_1} X_2}_{n-1}$$

is a weak equivalence.

The conditions for a P –Segal map to be a weak equivalence can be rephrased in terms of derived mapping spaces by means of the following notation.


Notation 1.16 Let P be a polygonal decomposition of the regular $(n+1)$ –gon, consisting of $(k+1)$ –gons of the form $\{i_0, \dots, i_k\}$ for varying k . We denote by $\Delta[P]$ the smallest subcomplex of $\Delta[n]$ containing all these simplices, as in [11, Formula 2.2.12]. The simplicial set $\Delta[P]$ can be expressed as a colimit of standard simplices

$$\operatorname{colim}_{\{i_0, \dots, i_k\} \in P} \Delta[k] \cong \Delta[P] \subset \Delta[n].$$

We emphasize two special cases. When $P = T$ is a triangulation of the regular $(n+1)$ –gon with vertices from the original $(n+1)$ –gon, the simplicial set $\Delta[T]$ can be expressed as an iterated pushout of the form

$$\underbrace{\Delta[2] \amalg_{\Delta[1]} \cdots \amalg_{\Delta[1]} \Delta[2]}_{n-1} \cong \Delta[P] \subset \Delta[n].$$

When P is a decomposition of the $(n+1)$ –gon into a triangle $\{0, 1, 2\}$ and the remaining n –gon, or into the triangle $\{n-2, n-1, n\}$ and the remaining n –gon, such as in the examples



the simplicial set $\Delta[P]$ can be expressed as a pushout of the form

$$\Delta[n-1] \amalg_{\Delta[1]} \Delta[2] \cong \Delta[P] \subset \Delta[n].$$

This discussion can be summarized via the following proposition:

Proposition 1.17 *Given a simplicial object X in \mathcal{C} , the P –Segal map is a weak equivalence in \mathcal{C} if and only if the associated map*

$$\mathrm{Map}^h(\Delta[n], X) \rightarrow \mathrm{Map}^h(\Delta[P], X),$$

induced by the inclusion

$$\mathrm{colim}_{\{i_0, \dots, i_k\} \in P} \Delta[k] \cong \Delta[P] \hookrightarrow \Delta[n]$$

is a weak equivalence in \mathcal{C} .

Determining whether a simplicial object is 2–Segal from the definition can be difficult, since there are so many possible triangulations of a polygon. However, our next result gives a substantial simplification, namely, that it is enough to check specific P –Segal maps induced by decompositions of an $(n+1)$ –gon into a triangle and remaining n –gon.

To state that result, we recall some notation from [11]. Given a simplicial object X in \mathcal{C} and an injective map $\theta: [k] \rightarrow [n]$ in Δ , we denote the induced map θ^* by $X_n \rightarrow X_{\{\theta(0), \dots, \theta(k)\}}$. Note that this notation is compatible with composition.

Proposition 1.18 *A simplicial object X in \mathcal{C} is 2–Segal if and only if, for any $n \geq 3$, the induced maps*

$$X_n \rightarrow X_{\{0,1,2\}} \overset{h}{\times}_{X_{\{0,2\}}} X_{\{0,2,\dots,n\}} \quad \text{and} \quad X_n \rightarrow X_{\{n-2,n-1,n\}} \overset{h}{\times}_{X_{\{n-2,n\}}} X_{\{0,\dots,n-2,n\}}$$

are weak equivalences.

Since we are working with model categories, it is convenient to make use of the following fact in the proof, allowing us to use strict pullbacks, rather than homotopy pullbacks, in the targets of the Segal maps.

Lemma 1.19 *If X is an injectively fibrant simplicial object in \mathcal{C} , then, for any injective map $\theta: [k] \rightarrow [n]$ in Δ , the induced map*

$$X_n \rightarrow X_{\{\theta(0), \dots, \theta(k)\}}$$

is a fibration in \mathcal{C} . In particular,

$$X_{\{0, \dots, j\}} \times_{X_{\{0, j\}}} X_{\{0, j, \dots, n\}} \simeq X_{\{0, \dots, j\}} \overset{h}{\times}_{X_{\{0, j\}}} X_{\{0, j, \dots, n\}}.$$

Moreover, for every $0 \leq j \leq n$, the projections

$$X_{\{0, \dots, j\}} \times_{X_{\{0, j\}}} X_{\{0, j, \dots, n\}} \rightarrow X_{\{0, \dots, j\}} \text{ and } X_{\{0, \dots, j\}} \times_{X_{\{0, j\}}} X_{\{0, j, \dots, n\}} \rightarrow X_{\{0, j, \dots, n\}}$$

are fibrations in \mathcal{C} .

Proof By Lemma 1.6, the map $\theta: \Delta[k] \rightarrow \Delta[n]$ is a levelwise cofibration between discrete simplicial objects. Since X is injectively fibrant, it follows that the induced map

$$X_n \cong \text{Map}(\Delta[n], X) \rightarrow \text{Map}(\Delta[k], X) \cong X_{\{\theta(0), \dots, \theta(k)\}}$$

is a fibration in \mathcal{C} , as desired. □

We can now prove our criterion for 2–Segal objects.

Proof of Proposition 1.18 In [11, Proposition 2.3.2], Dyckerhoff and Kapranov show that X is 2–Segal if and only if, for every $n \geq 3$, $0 < i < n - 1$ and $1 < j < n$, the maps

$$X_n \rightarrow X_{\{0, \dots, j\}} \overset{h}{\times}_{X_{\{0, j\}}} X_{\{0, j, \dots, n\}} \quad \text{and} \quad X_n \rightarrow X_{\{i, \dots, n\}} \overset{h}{\times}_{X_{\{i, n\}}} X_{\{0, \dots, i, n\}}$$


are weak equivalences. With $j = 2$ and $i = n - 2$, we get the direct implication. For the converse implication, we proceed by induction on n . Here we show that, for every n and every $1 < j < n$, the map

$$(1.20) \quad X_n \rightarrow X_{\{0, \dots, j\}} \overset{h}{\times}_{X_{\{0, j\}}} X_{\{0, j, \dots, n\}}$$

is a weak equivalence; the argument for the other map follows by a similar argument.

When $n = 3$, we have that $j = 2$, so the map in (1.20) is by assumption a weak equivalence. Fix $n > 3$ and assume the condition holds for $k < n$. For $1 < j < n$,

consider the decomposition of the $(n+1)$ –gon given by the two diagonals $\{0, j\}$ and $\{0, 2\}$,



into a triangle and two other polygons. By the inductive hypothesis, the maps

$$\begin{aligned} X_{\{0,\dots,j\}} &\rightarrow X_{\{0,1,2\}} \overset{h}{\times}_{X_{\{0,2\}}} X_{\{0,2,\dots,j\}}, \\ X_{\{0,2,\dots,n\}} &\rightarrow X_{\{0,2,\dots,j\}} \overset{h}{\times}_{X_{\{0,j\}}} X_{\{0,j,\dots,n\}} \end{aligned}$$

are weak equivalences.

We exhibit the desired map as a zigzag of weak equivalences using the two-out-of-three property, but to get the commuting diagram, it is useful to use concrete models for the homotopy pullback. We do so by first taking an injective fibrant replacement for X . Using [Lemma 1.19](#) we can now work with strict pullbacks rather than homotopy pullbacks. Thus, in the diagram

$$\begin{array}{ccc} X_n & \longrightarrow & X_{\{0,\dots,j\}} \times_{X_{\{0,j\}}} X_{\{0,j,\dots,n\}} \\ \downarrow \simeq & & \downarrow \simeq \\ X_{\{0,1,2\}} \times_{X_{\{0,2\}}} X_{\{0,2,\dots,n\}} & \xrightarrow{\simeq} & X_{\{0,1,2\}} \times_{X_{\{0,2\}}} X_{\{0,2,\dots,j\}} \times_{X_{\{0,j\}}} X_{\{0,j,\dots,n\}} \end{array}$$

the bottom and right arrows are weak equivalences. The left arrow is a weak equivalence by assumption. The two-out-of-three property implies that the top arrow is a weak equivalence as well, as we wished to show. \square

Remark 1.21 There is a technically subtle step in the proof above which arises from working with homotopy pullbacks in the model category itself rather than in the corresponding $(\infty, 1)$ –category or homotopy category. There are different ways to handle this subtlety; we chose [Lemma 1.19](#) as the most suitable for our purposes. Another possibility is to choose a functorial model for homotopy pullbacks in \mathcal{C} . We continue to use the proof strategy as above in future proofs without further discussion.

In both [\[11\]](#) and [\[16\]](#), the following unitality condition was shown to be satisfied for a number of examples. Roughly speaking, it says that identity maps behave as we expect them to.

Definition 1.22 A 2–Segal object in \mathcal{C} is *unital* if, for all $n \geq 2$ and $0 \leq i \leq n - 1$, the diagram

$$\begin{array}{ccc} X_{n-1} & \xrightarrow{\alpha_i} & X_0 \\ s_i \downarrow & & \downarrow s_0 \\ X_n & \xrightarrow{\beta_i} & X_1 \end{array}$$

is a homotopy pullback, where s_0 and s_i are degeneracy maps and the maps α_i and β_i are induced by the maps

$$\alpha^i : [0] \rightarrow [n - 1], \quad 0 \mapsto i, \quad \text{and} \quad \beta^i : [1] \rightarrow [n], \quad 0 < 1 \mapsto i < i + 1.$$

Remark 1.23 The squares appearing in the definition of unitality are induced by the square of simplicial sets

$$\begin{array}{ccc} \Delta[n - 1] & \xleftarrow{\alpha^i} & \Delta[0] \\ s^i \uparrow & & \uparrow s^0 \\ \Delta[n] & \xleftarrow{\beta^i} & \Delta[1] \end{array}$$

As this square is commutative, it induces a map

(1.24)
$$\Delta[n] \amalg_{\Delta[1]}^{\beta^i, s^0} \Delta[0] \rightarrow \Delta[n - 1].$$

We can now rephrase the definition of unitality by saying that X is a unital 2–Segal object if and only if all induced maps

$$\mathrm{Map}^h(\Delta[n - 1], X) \rightarrow \mathrm{Map}^h(\Delta[n] \amalg_{\Delta[1]}^{\beta^i, s^0} \Delta[0], X)$$

are weak equivalences.

However, Feller, Garner, Kock, Underhill-Proulx and Weber recently showed that the unitality condition is automatically satisfied for any 2–Segal object:

Theorem 1.25 [12] *Every 2–Segal object is unital.*

In particular, we do not need to establish separately that our 2–Segal objects of interest are unital. However, this condition still plays an important role in proving some of our comparisons of model structures, so we cannot ignore it entirely.

Remark 1.26 For the rest of this paper, we typically omit the term “unital” when referring to 2–Segal objects, with two main exceptions. First, when we give formal statements of prior results which use the term, we include it for clarity and consistency. Second, as mentioned above, we occasionally need to invoke the fact that 2–Segal objects are indeed unital.

1.3 2–Segal sets and augmented stable double categories

In our previous paper [6], we established an equivalence of categories between 2–Segal sets and certain kinds of double categories. In this subsection, we review double categories, the extra conditions we want to place on them, and the equivalence with 2–Segal sets.

In brief, a *double category* \mathcal{D} is a category internal to categories. In other words, it consists of objects $\text{Ob } \mathcal{D}$, horizontal morphisms $\text{Hor } \mathcal{D}$, vertical morphisms $\text{Ver } \mathcal{D}$ and squares $\text{Sq } \mathcal{D}$. We denote horizontal morphisms by \rightrightarrows and vertical morphisms by \rightrightarrows . There are usual source and target maps $s_h, t_h: \text{Hor } \mathcal{D} \rightarrow \text{Ob } \mathcal{D}$ and $s_v, t_v: \text{Ver } \mathcal{D} \rightarrow \text{Ob } \mathcal{D}$. There are “horizontal” source and target maps $s_h, t_h: \text{Sq } \mathcal{D} \rightarrow \text{Ver } \mathcal{D}$ and “vertical”¹ source and target maps $s_v, t_v: \text{Sq } \mathcal{D} \rightarrow \text{Hor } \mathcal{D}$. We refer the reader to [6, Section 3] for more details on the notation that we use, and to [14] or [17] for the general theory of double categories.

We recall from [6] that a double category \mathcal{D} is *stable* if the maps

$$(1.27) \quad \text{Hor } \mathcal{D} \times_{\text{Ob } \mathcal{D}}^{s_v} \text{Ver } \mathcal{D} \xleftarrow{(s_v, s_h)} \text{Sq } \mathcal{D} \xrightarrow{(t_h, t_v)} \text{Ver } \mathcal{D} \times_{\text{Ob } \mathcal{D}}^{t_v} \text{Hor } \mathcal{D}$$

are isomorphisms. These isomorphisms amount to requiring that every span in \mathcal{D} given by a horizontal arrow and a vertical arrow can be filled uniquely to a square, and similarly for every cospan.

A double category \mathcal{D} is *augmented* by a subset $A\mathcal{D}$ of its objects if the maps

$$(1.28) \quad A\mathcal{D} \times_{\text{Ob } \mathcal{D}}^{s_h} \text{Hor } \mathcal{D} \xrightarrow{t_h \circ \text{pr}_2} \text{Ob } \mathcal{D} \quad \text{and} \quad \text{Ver } \mathcal{D} \times_{\text{Ob } \mathcal{D}}^{t_v} A\mathcal{D} \xrightarrow{s_v \circ \text{pr}_1} \text{Ob } \mathcal{D}$$

are isomorphisms. These isomorphisms amount to requiring that for every object x in \mathcal{D} there is a unique horizontal morphism with source in the augmentation and target x , and, similarly, there is a unique vertical morphism with source x and target in the augmentation.

We recall three important examples of stable double categories that appear later in the paper.

Example 1.29 For any $n \geq 0$, consider the double category $\mathcal{W}[n]$ of [6, Definition 4.1] defined as follows:

- The set of objects is given by

$$\text{Ob}(\mathcal{W}[n]) = \{(i, j) \mid 0 \leq i \leq j \leq n\}.$$

¹Although possibly confusing, the “horizontal” source and target correspond to horizontal composition, and hence are vertical morphisms.

- The set of horizontal arrows is given by

$$\text{Hor}(\mathcal{W}[n]) = \{(i, j, \ell) \mid 0 \leq i \leq j \leq \ell \leq n\}.$$

A horizontal morphism (i, j, ℓ) with $i \leq j \leq \ell$ is viewed as a map

$$(i, j) \twoheadrightarrow (i, \ell).$$

- The set of vertical arrows is given by

$$\text{Ver}(\mathcal{W}[n]) = \{(i, k, j) \mid 0 \leq i \leq k \leq j \leq n\}.$$

A vertical morphism (i, k, j) with $i \leq k \leq j$ is viewed as a map

$$(i, j) \twoheadrightarrow (k, j).$$

- The set of squares is given by

$$\text{Sq}(\mathcal{W}[n]) = \{(i, k, j, \ell) \mid 0 \leq i \leq k \leq j \leq \ell \leq n\}.$$

A square (i, k, j, ℓ) with $0 \leq i \leq k \leq j \leq \ell \leq n$ is viewed as a square

(1.30)

$$\begin{array}{ccc} (i, j) & \twoheadrightarrow & (i, \ell) \\ \downarrow & \searrow & \downarrow \\ (k, j) & \twoheadrightarrow & (k, \ell) \end{array}$$

For any $n \geq 0$ the double category $\mathcal{W}[n]$ is stable, and augmented when endowed with the augmentation set

$$A(\mathcal{W}[n]) = \{(i, i) \mid 0 \leq i \leq n\}.$$

For instance, if we follow the convention that a diagram represents a double category that contains all the displayed objects, arrows and squares, and all their possible composites and identities, and that stars $*$ denote elements of the augmentation set, the double category $\mathcal{W}[4]$ can be depicted as follows:

(1.31)

Example 1.32 For any $n \geq 0$, consider the sub-double category $\mathcal{H}[n]$ of $\mathcal{W}[n]$ spanned by the first row of $\mathcal{W}[n]$, ie the full sub-double category on the set

$$\text{Ob}(\mathcal{H}[n]) = \{(0, j) \mid 0 \leq j \leq n\} \cong \{j \mid 0 \leq j \leq n\}.$$

For any n the double category $\mathcal{H}[n]$ is stable, but it cannot be augmented by any subset of its objects.

For example, the double category $\mathcal{H}[4]$ can be depicted as follows:

$$(1.33) \quad * \rightharpoonup \bullet \rightharpoonup \bullet \rightharpoonup \bullet \rightharpoonup \bullet$$

Note that $\mathcal{H}[n]$ only has identity squares. Dually, we can consider the double category $\mathcal{V}[n]$ spanned by the last column of $\mathcal{W}[n]$, which is also stable but not augmented.

Example 1.34 For any $q, r \geq 0$, consider the double category $[q] \boxtimes [r]$ from [14, Example 2.9], defined as follows:

- The set of objects is given by

$$\text{Ob}([q] \boxtimes [r]) = \{(i, j) \mid 1 \leq i \leq q, 1 \leq j \leq r\}.$$

- The set of horizontal arrows is given by

$$\text{Hor}([q] \boxtimes [r]) = \{(i, j, k) \mid 0 \leq i \leq q, 0 \leq j \leq k \leq r\}.$$

- The set of vertical arrows is given by

$$\text{Ver}([q] \boxtimes [r]) = \{(i, j, k) \mid 0 \leq i \leq k \leq q, 0 \leq j \leq r\}.$$

- The set of squares is given by

$$\text{Sq}([q] \boxtimes [r]) = \{(i, j, k, \ell) \mid 0 \leq i \leq k \leq q, 0 \leq j \leq \ell \leq r\}.$$

For instance, the double category $[2] \boxtimes [3]$ can be depicted as follows:

$$(1.35) \quad \begin{array}{ccccccc} \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet \\ \downarrow & \searrow & \downarrow & \searrow & \downarrow & \searrow & \downarrow \\ \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet \\ \downarrow & \searrow & \downarrow & \searrow & \downarrow & \searrow & \downarrow \\ \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet \end{array}$$

For any q and r , the double category $[q] \boxtimes [r]$ is stable. However, it cannot be made into an augmented double category, since the sets of connected components of the horizontal category and of the vertical category are necessarily in bijection for an augmented double category.

We denote by $\mathcal{DCat}_{\text{aug}}^{\text{st}}$ the category of augmented stable double categories, and by $\mathcal{U}2\text{Seg}$ the category of unital 2–Segal sets. Note that we include “unital” here for emphasis to be consistent with the earlier work which we reference here. In our previous paper [6], we defined the functors

$$\mathcal{P}: \mathcal{U}2\text{Seg} \rightarrow \mathcal{DCat}_{\text{aug}}^{\text{st}} \quad \text{and} \quad S_{\bullet}: \mathcal{DCat}_{\text{aug}}^{\text{st}} \rightarrow \mathcal{U}2\text{Seg},$$

which we briefly recall. For more details, see [6, Sections 4 and 5].

Given a unital 2–Segal set X , the *path construction* \mathcal{P} assigns to X an augmented stable double category $\mathcal{P}X$ in which $\text{Ob}(\mathcal{P}X) = X_1$, $\text{Hor}(\mathcal{P}X) = X_2 = \text{Ver}(\mathcal{P}X)$, $\text{Sq}(\mathcal{P}X) = X_3$ and $A(\mathcal{P}X) = X_0$. The source, target and identity maps are given by appropriate face and degeneracy maps, and the various composites are induced by the inverses of the 2–Segal maps.

In the other direction, given an augmented stable double category \mathcal{D} , the *Waldhausen construction* S_{\bullet} takes \mathcal{D} to a simplicial set whose set of n –simplices is corepresented by $\mathcal{W}[n]$.

Remark 1.36 Although Waldhausen did not work explicitly with double categories in [32], his S_{\bullet} –construction can be interpreted in that language. Given a Waldhausen category \mathcal{F} , one can define an augmented (not necessarily stable) double category $\mathcal{D}(\mathcal{F})$ with cofibrations, quotient maps, pushout diagrams and a chosen zero object as the horizontal morphisms, vertical morphisms, squares and the augmentation, respectively. Then the objects of $S_{\bullet}\mathcal{F}$ can be identified with the set of augmented double functors $\mathcal{W}[n] \rightarrow \mathcal{D}(\mathcal{F})$. We revisit this point of view in [7].

The following theorem is the main result of [6]:

Theorem 1.37 [6, Theorem 6.1] *The path construction \mathcal{P} and the Waldhausen construction S_{\bullet} induce an equivalence of categories*

$$\mathcal{P}: \mathcal{U}2\text{Seg} \rightleftarrows \mathcal{DCat}_{\text{aug}}^{\text{st}} : S_{\bullet}.$$

The aim of this paper is to prove the homotopical analogue to this theorem. Before moving away from the discrete setting, however, we establish a helpful result about the effect of applying the path construction to a simplex.

For any $n \geq 0$, the simplicial set $\Delta[n]$ is the nerve of the category $[n]$ and hence a Segal set. We can apply the functor \mathcal{P} to it (since it is automatically unital) to obtain an augmented stable double category.

Proposition 1.38 *There is a natural isomorphism of augmented stable double categories*

$$\mathcal{P}\Delta[n] \cong \mathcal{W}[n].$$

Proof By the construction of \mathcal{P} from [6, Section 5], we have the following descriptions of the sets of objects, horizontal and vertical morphisms, squares and augmentation set of the augmented stable double category $\mathcal{P}\Delta[n]$:

$$\mathrm{Ob}(\mathcal{P}\Delta[n]) = \Delta[n]_1 \cong \mathrm{Hom}([1], [n]) \cong \{(i, j) \mid 0 \leq i \leq j \leq n\},$$

$$\mathrm{Hor}(\mathcal{P}\Delta[n]) = \Delta[n]_2 \cong \mathrm{Hom}([2], [n]) \cong \{(i, j, \ell) \mid 0 \leq i \leq j \leq \ell \leq n\},$$

$$\mathrm{Ver}(\mathcal{P}\Delta[n]) = \Delta[n]_2 \cong \mathrm{Hom}([2], [n]) \cong \{(i, k, j) \mid 0 \leq i \leq k \leq j \leq n\},$$

$$\mathrm{Sq}(\mathcal{P}\Delta[n]) = \Delta[n]_3 \cong \mathrm{Hom}([3], [n]) \cong \{(i, k, j, \ell) \mid 0 \leq i \leq k \leq j \leq \ell \leq n\},$$

$$A(\mathcal{P}\Delta[n]) = \Delta[n]_0 \cong \mathrm{Hom}([0], [n]) \cong \{(i, i) \mid 0 \leq i \leq n\}.$$

It remains to compare the rest of the double category structure: the horizontal and vertical source and target maps, identities and compositions.

By definition, the horizontal source and target maps $\mathrm{Hor}(\mathcal{P}\Delta[n]) \rightrightarrows \mathrm{Ob}(\mathcal{P}\Delta[n])$ are given by $d_2, d_1: \Delta[n]_2 \rightrightarrows \Delta[n]_1$, respectively, and therefore send an element $(i, j, \ell) \in \mathrm{Hor}(\mathcal{P}\Delta[n])$ to (i, j) and (i, ℓ) , which are exactly the source and target of the horizontal morphism $(i, j, \ell): (i, j) \rightarrowtail (i, \ell)$ in $\mathcal{W}[n]$.

Similarly, the vertical source and target maps $\mathrm{Ver}(\mathcal{P}\Delta[n]) \rightrightarrows \mathrm{Ob}(\mathcal{P}\Delta[n])$ are given by $d_1, d_0: \Delta[n]_2 \rightrightarrows \Delta[n]_1$, which coincide with the source and target maps of the vertical morphism $(i, k, j): (i, j) \rightarrowtail (k, j)$ in $\mathcal{W}[n]$.

On $\mathrm{Sq}(\mathcal{P}\Delta[n]) = \Delta[n]_3$, the vertical source and target by definition are given by $d_1, d_0: \Delta[n]_3 \rightrightarrows \Delta[n]_2$, whereas the horizontal source and target are given by

$$d_3, d_2: \Delta[n]_3 \rightrightarrows \Delta[n]_2.$$

By inspection, for an element $(i, k, j, \ell) \in \mathrm{Sq}(\mathcal{P}\Delta[n])$, these definitions coincide with those indicated in (1.30).

A straightforward check establishes that composition and identities of $\mathcal{P}\Delta[n]$ and $\mathcal{W}[n]$ coincide. \square

2 The generalized S_\bullet -construction

In this section we introduce our main construction, a homotopical generalization of the S_\bullet -construction as described in the previous section.

The first step is to establish the correct input objects for this construction, which should possess some of the features of an exact category. As we saw in the previous section, in the discrete case the appropriate structure is that of an augmented stable double category. Here, we generalize double categories to double Segal objects, then develop augmented and stable versions in this context.

2.1 Double Segal objects

Inspired by the fact that double categories are categories internal to categories, we use a higher categorical version thereof modeled by *double Segal objects in \mathcal{C}* , as investigated by Haugseng [19]. They are Segal objects in the category of Segal objects in \mathcal{C} and model double categories up to homotopy.

It is important to note that double Segal objects are more general than the 2-fold Segal spaces of [2] or [26], which instead model $(\infty, 2)$ -categories, or 2-categories up to homotopy.

Let us look at this definition more precisely. Recall the notion of the spine of a standard simplex from Notation 1.12.

Definition 2.1 A bisimplicial object Y in \mathcal{C} is a *double Segal object* if for every $q, r \geq 1$ the maps

$$Y_{q,r} \rightarrow \underbrace{Y_{q,1} \overset{h}{\times}_{Y_{q,0}} \cdots \overset{h}{\times}_{Y_{q,0}} Y_{q,1}}_r \quad \text{and} \quad Y_{q,r} \rightarrow \underbrace{Y_{1,r} \overset{h}{\times}_{Y_{0,r}} \cdots \overset{h}{\times}_{Y_{0,r}} Y_{1,r}}_q$$

are weak equivalences in \mathcal{C} . Here the left-hand map is induced by the inclusion $\mathcal{I}[r] \hookrightarrow \Delta[r]$ in the second variable in $(\Delta \times \Delta)^{\text{op}}$, whereas the right-hand map is induced by the same inclusion $\mathcal{I}[q] \hookrightarrow \Delta[q]$, but in the first variable.

Let us consider two motivating examples.

Example 2.2 When \mathcal{C} is the category of sets with its trivial model structure, a double Segal set Y is precisely an appropriately defined nerve of a double category \mathcal{D} , where $Y_{0,0} = \text{Ob } \mathcal{D}$ is the set of objects, $Y_{0,1} = \text{Hor } \mathcal{D}$ and $Y_{1,0} = \text{Ver } \mathcal{D}$ are the sets of horizontal and vertical morphisms, respectively, and $Y_{1,1} = \text{Sq } \mathcal{D}$ is the set of distinguished squares [14, Definition 5.12]. More generally, using Example 1.34, the set $Y_{q,r} = \text{Hom}_{\mathcal{D}\text{Cat}}([q] \boxtimes [r], \mathcal{D})$ is the set of $q \times r$ grids of distinguished squares.

Horizontal composition of squares is a strict operation encoded by

$$Y_{1,1} \times_{Y_{1,0}} Y_{1,1} \xleftarrow{\cong} Y_{1,2} \rightarrow Y_{1,1}.$$

The rest of the structure can be defined similarly.

Example 2.3 Consider the case when \mathcal{C} is the category of simplicial sets with the classical model structure due to Quillen [28]. Just as Segal spaces can be regarded as a homotopical analogue of Segal sets, which are nerves of categories, double Segal spaces give a homotopical version of double Segal sets, which we have just seen model nerves of double categories.


Given a double Segal space Y , we can think of $Y_{0,0}$ as a space of objects, $Y_{0,1}$ and $Y_{1,0}$ as spaces of horizontal and vertical morphisms, respectively, and $Y_{1,1}$ as a space of distinguished squares. Horizontal composition of squares is an operation now only well defined up to homotopy, and encoded by the span

$$Y_{1,1} \mathop{\times}\limits^h_{Y_{1,0}} Y_{1,1} \xleftarrow{\cong} Y_{1,2} \rightarrow Y_{1,1},$$

where the homotopy fiber product on the left-hand side encodes the space of horizontally composable squares, and $Y_{1,2}$ represents the space of horizontally composable squares together with a specified composition of such. According to this interpretation, $Y_{q,r}$ can be thought of as the space of $q \times r$ grids of distinguished squares together with their intermediate composites.

Remark 2.4 An important feature of a double category is the *interchange law*, which describes the compatibility of horizontal and vertical composition. Let us look more closely at how the analogous law for a double Segal object is encoded by the bisimplicial structure.

Consider the space \mathbb{Y} of four adjacent squares in a double Segal space Y (but without any composition) glued along the dots as in the left picture below. In contrast, the picture on the right is supposed to possess all composites of squares, with appropriate compatibility, and corresponds to an element in $Y_{2,2}$:



More precisely, by “glued along the dots” we mean that the space \mathbb{Y} can be expressed as the homotopy limit of a diagram of the form

$$\mathbb{Y} = \left(\begin{array}{ccccc} Y_{1,1} & \rightarrow & Y_{1,0} & \leftarrow & Y_{1,1} \\ \downarrow & & \downarrow & & \downarrow \\ Y_{0,1} & \rightarrow & Y_{0,0} & \leftarrow & Y_{0,1} \\ \uparrow & & \uparrow & & \uparrow \\ Y_{1,1} & \rightarrow & Y_{1,0} & \leftarrow & Y_{1,1} \end{array} \right),$$

which in turn can be written as either of the iterated homotopy fiber products

$$\begin{aligned} &(Y_{1,1} \overset{h}{\times}_{Y_{0,1}} Y_{1,1}) \overset{h}{\times}_{(Y_{1,0} \overset{h}{\times}_{Y_{0,0}} Y_{1,0})} (Y_{1,1} \overset{h}{\times}_{Y_{0,1}} Y_{1,1}), \\ &(Y_{1,1} \overset{h}{\times}_{Y_{1,0}} Y_{1,1}) \overset{h}{\times}_{(Y_{0,1} \overset{h}{\times}_{Y_{0,0}} Y_{0,1})} (Y_{1,1} \overset{h}{\times}_{Y_{1,0}} Y_{1,1}). \end{aligned}$$

Consider the following diagram, which commutes up to homotopy:

$$\begin{array}{ccccc} (Y_{1,1} \overset{h}{\times}_{Y_{0,1}} Y_{1,1}) \overset{h}{\times}_{(Y_{1,0} \overset{h}{\times}_{Y_{0,0}} Y_{1,0})} (Y_{1,1} \overset{h}{\times}_{Y_{0,1}} Y_{1,1}) & \xleftarrow{\simeq} & Y_{2,1} \overset{h}{\times}_{Y_{2,0}} Y_{2,1} & \rightarrow & Y_{1,1} \overset{h}{\times}_{Y_{1,0}} Y_{1,1} \\ \uparrow \simeq & & \uparrow \simeq & \nearrow & \simeq \uparrow \\ \text{holim } \mathbb{Y} & \xleftarrow{\hspace{2cm}} & Y_{2,2} & \rightarrow & Y_{1,2} \\ & & & \searrow & \downarrow \\ & & & & Y_{1,1} \\ \downarrow \simeq & & \downarrow \simeq & \nearrow & \uparrow \\ (Y_{1,1} \overset{h}{\times}_{Y_{1,0}} Y_{1,1}) \overset{h}{\times}_{(Y_{0,1} \overset{h}{\times}_{Y_{0,0}} Y_{0,1})} (Y_{1,1} \overset{h}{\times}_{Y_{1,0}} Y_{1,1}) & \xleftarrow{\simeq} & Y_{1,2} \overset{h}{\times}_{Y_{0,2}} Y_{1,2} & \rightarrow & Y_{1,1} \overset{h}{\times}_{Y_{0,1}} Y_{1,1} \\ & & & & \downarrow \simeq \\ & & & & Y_{2,1} \end{array}$$

Here, reaching $Y_{1,1}$ through the top line corresponds to first composing the two pairs of squares vertically, and then composing the result horizontally. On the other hand, the bottom line corresponds to first composing the two pairs of squares horizontally, and then composing the result vertically. The two operations are therefore compatible with each other, as expected. One can similarly express higher versions of the interchange law by using similar versions of this diagram.

2.2 Stable double Segal objects

We now introduce the stability condition on a double Segal object, which generalizes the definition given in the discrete case. It also appears in work of Carlier [10].

For ease of notation, we denote an object $([k],[\ell])$ in $\Delta \times \Delta$ simply by (k,ℓ) . We denote by $s^h, t^h \colon (k,0) \rightarrow (k,\ell)$ the maps given by (id, α^0) and (id, α^ℓ) , respectively,

where α^i is the map from [Definition 1.22](#). We similarly denote by $s^v, t^v: (0, \ell) \rightarrow (k, \ell)$ the maps (α^0, id) and (α^k, id) , respectively.

Definition 2.5 A bisimplicial object Y in \mathcal{C} is *stable* if for every $q, r \geq 1$ the squares

$$(2.6) \quad \begin{array}{ccc} (0, 0) & \xrightarrow{s^h} & (0, r) \\ s^v \downarrow & & \downarrow s^v \\ (q, 0) & \xrightarrow{s^h} & (q, r) \end{array} \quad \text{and} \quad \begin{array}{ccc} (0, 0) & \xrightarrow{t^h} & (0, r) \\ t^v \downarrow & & \downarrow t^v \\ (q, 0) & \xrightarrow{t^h} & (q, r) \end{array}$$

induce weak equivalences

$$Y_{0,r} \overset{h}{\times}_{Y_{0,0}} Y_{q,0} \xleftarrow{\cong} Y_{q,r} \xrightarrow{\cong} Y_{q,0} \overset{h}{\times}_{Y_{0,0}} Y_{0,r}.$$

Under the double Segal condition, there is an easier criterion to check whether a bisimplicial object is stable. For an analogous result, see [\[10, Lemma 2.3.3\]](#).

Lemma 2.7 A double Segal space Y is stable if and only if the squares

$$(2.8) \quad \begin{array}{ccc} (0, 0) & \xrightarrow{s^h} & (0, 1) \\ s^v \downarrow & & \downarrow s^v \\ (1, 0) & \xrightarrow{s^h} & (1, 1) \end{array} \quad \text{and} \quad \begin{array}{ccc} (0, 0) & \xrightarrow{t^h} & (0, 1) \\ t^v \downarrow & & \downarrow t^v \\ (1, 0) & \xrightarrow{t^h} & (1, 1) \end{array}$$

induce weak equivalences

$$Y_{0,1} \overset{h}{\times}_{Y_{0,0}} Y_{1,0} \xleftarrow{\cong} Y_{1,1} \xrightarrow{\cong} Y_{1,0} \overset{h}{\times}_{Y_{0,0}} Y_{0,1}.$$

Proof We prove that for a double Segal object, the left diagram in condition [\(2.8\)](#) is equivalent to the left diagram in condition [\(2.6\)](#), and the equivalence of the right diagrams follows similarly. We proceed by induction on $q + r$. In the base case $q = r = 1$, the claim is exactly the assumption. Now, if $q + r > 2$, either $q > 1$ or $r > 1$. Without loss of generality, assume $q > 1$. We want to show that the left vertical map in the following homotopy commutative diagram (see [Remark 1.21](#)) is an equivalence:


$$\begin{array}{ccccc} Y_{q,r} & \xrightarrow{\cong} & Y_{1,r} \overset{h}{\times}_{Y_{0,r}} Y_{q-1,r} & \xrightarrow{\cong} & Y_{1,r} \overset{h}{\times}_{Y_{0,r}} (Y_{0,r} \overset{h}{\times}_{Y_{0,0}} Y_{q-1,0}) \\ \downarrow & & & & \downarrow \cong \\ & & & & (Y_{1,r} \overset{h}{\times}_{Y_{0,r}} Y_{0,r}) \overset{h}{\times}_{Y_{0,0}} Y_{q-1,0} \\ & & & & \downarrow \cong \\ Y_{0,r} \overset{h}{\times}_{Y_{0,0}} Y_{q,0} & \xrightarrow{\cong} & Y_{0,r} \overset{h}{\times}_{Y_{0,0}} Y_{1,0} \overset{h}{\times}_{Y_{0,0}} Y_{q-1,0} & \xleftarrow{\cong} & Y_{1,r} \overset{h}{\times}_{Y_{0,0}} Y_{q-1,0} \end{array}$$

The left-hand horizontal maps are given by Segal maps in the first index, and are equivalences since Y is double Segal. The right-hand horizontal maps are equivalences by the induction hypothesis. The vertical left maps use the properties of the homotopy fiber product. The two-out-of-three property finishes the proof. \square

2.3 Preaugmented bisimplicial objects

Recall from Section 1.3 that in the discrete context, we singled out some collection of the objects of a double category which in turn satisfied some conditions. In this section, we focus on identifying such a collection for a more general double Segal object. Since we do not yet impose any universality conditions, we refer to such objects as *preaugmented*. We do so by modifying the indexing category $\Delta \times \Delta$.

Definition 2.9 Let Σ be the category obtained from $\Delta \times \Delta$ by adding a new terminal object, denoted by $[-1]$:



A *preaugmented bisimplicial object* in \mathcal{C} is a functor $Y: \Sigma^{\text{op}} \rightarrow \mathcal{C}$. The category of preaugmented bisimplicial objects is the category $\text{Fun}(\Sigma^{\text{op}}, \mathcal{C})$, which, for simplicity of notation, we henceforth denote by $\mathcal{C}^{\Sigma^{\text{op}}}$. We denote other categories of functors similarly; for example, we write $\text{Fun}((\Delta \times \Delta)^{\text{op}}, \mathcal{C})$ as $\mathcal{C}^{(\Delta \times \Delta)^{\text{op}}}$.

Definition 2.10 Let Y be a preaugmented bisimplicial object in \mathcal{C} . Its *underlying bisimplicial object* is the image of Y under the functor

$$i^*: \mathcal{C}^{\Sigma^{\text{op}}} \rightarrow \mathcal{C}^{(\Delta \times \Delta)^{\text{op}}}$$

induced by the inclusion $i: \Delta \times \Delta \rightarrow \Sigma$.

Remark 2.11 The category Σ can be expressed as a pushout of categories

$$\Sigma \cong (\Delta \times \Delta) \amalg_{[0]}^{d^1} [1].$$

Therefore, the category of preaugmented bisimplicial objects can be expressed as a pullback of categories

$$\mathcal{C}^{\Sigma^{\text{op}}} \cong \mathcal{C}^{(\Delta \times \Delta)^{\text{op}}} \times_{\mathcal{C}} \mathcal{C}^{[1]^{\text{op}}},$$

decomposing the data of a preaugmented bisimplicial object into its bisimplicial part and its preaugmentation.

Let us consider some examples when \mathcal{C} is the category of sets; via [Proposition 1.1](#) we can consider these examples as discrete objects for more general \mathcal{C} . We begin with representable functors.

Example 2.12 Let \mathcal{C} be the category of sets. For any $q, r \geq 0$ we have the representable preaugmented bisimplicial set $\Sigma[q, r]$ given by

$$\Sigma[q, r]_{k, \ell} = \text{Hom}_{\Sigma}((k, \ell), (q, r)) \cong \text{Hom}_{\Delta \times \Delta}((k, \ell), (q, r))$$

for every $k, \ell \geq 0$, and

$$\Sigma[q, r]_{-1} = \text{Hom}_{\Sigma}([-1], (q, r)) = \emptyset.$$

Note that the bisimplicial set $i^* \Sigma[q, r]$ (without the preaugmentation) is precisely the double nerve of the double category $[q] \boxtimes [r]$ from [Example 1.34](#). Thus, one can keep the picture (1.35) in mind to visualize $\Sigma[q, r]$. These ideas are discussed in more detail in [Section 3.1](#).

However, we must treat the functor represented by $[-1]$ separately. Since $[-1]$ is the terminal object of Σ , necessarily $\Sigma[-1]$ is the constant functor at $\{*\}$.

The next examples are motivated by [Examples 1.32](#) and [1.29](#); a more explicit connection will be established in the next section.

Example 2.13 Again, let \mathcal{C} be the category of sets. For any $n \geq 0$, we define preaugmented bisimplicial sets

$$H[n] := \Sigma[0, n]^{s^h} \amalg_{\Sigma[0, 0]} \Sigma[-1] \quad \text{and} \quad V[n] := \Sigma[n, 0]^{t^v} \amalg_{\Sigma[0, 0]} \Sigma[-1],$$

where $s^h: \Sigma[0, 0] \rightarrow \Sigma[0, n]$ is induced by composition with s^h in $\Delta \times \Delta$ and

$$\Sigma[0, 0]_{k, \ell} \cong \text{Hom}_{\Delta \times \Delta}((k, \ell), (0, 0)) \xrightarrow{s^h \circ -} \text{Hom}_{\Delta \times \Delta}((k, \ell), (0, n)) \cong \Sigma[0, n]_{k, \ell}.$$

Similarly, $\iota^v: \Sigma[0, 0] \rightarrow \Sigma[n, 0]$ is induced by composition with ι^v in $\Delta \times \Delta$.

In particular, $H[0] = \Sigma[-1] = V[0]$. In the next section, after defining a nerve functor for augmented stable double categories, we recover $H[n]$ as the nerve of the double category $\mathcal{H}[n]$ from [Example 1.32](#), and analogously for $V[n]$. Nonetheless, one can refer to [\(1.33\)](#) for an intuitive description for $H[n]$.

Finally, we consider an analogue of [Example 1.29](#).

Example 2.14 For any $n \geq 0$, the preaugmented bisimplicial set $W[n]$ is defined by

$$W[n]_{k,\ell} = \{(i_0, \dots, i_k, j_0, \dots, j_\ell) \mid 0 \leq i_0 \leq \dots \leq i_k \leq j_0 \leq \dots \leq j_\ell \leq n\}$$

for any $k, \ell \geq 0$ and by

$$W[n]_{-1} = \{(i, i) \mid 0 \leq i \leq n\}.$$

Degeneracy maps are given by repeating the appropriate index, face maps by removing the designated index, and the augmentation map by the canonical inclusion. It is useful to note that $W[0] = \Sigma[-1]$.

Also in this case, after defining the nerve functor, we can recover $W[n]$ as the nerve of $\mathcal{W}[n]$. In the meantime, the reader may keep the diagram [\(1.31\)](#) in mind for intuition for $W[n]$.

The three examples $H[n]$, $V[n]$ and $W[n]$, taken as discrete objects in the setting of more general simplicial objects in \mathcal{C} , play an important role in what follows.

2.4 The generalized path construction and the S_\bullet -construction

While we still need to introduce the notion of augmentation in this context, the structure of a preaugmented simplicial object is sufficient to define the functors which we use for our comparison with 2–Segal objects. In this section we introduce an adjoint pair of functors between the category of simplicial objects in \mathcal{C} and the category of preaugmented bisimplicial objects in \mathcal{C} . These functors generalize in a precise sense the path construction and the S_\bullet -construction from [Theorem 1.37](#), as we show explicitly in [Proposition 3.6](#).

Definition 2.15 The ordinal sum $\Delta \times \Delta \rightarrow \Delta$ extends to a functor $p: \Sigma \rightarrow \Delta$ along the canonical inclusion $i: \Delta \times \Delta \rightarrow \Sigma$ satisfying

$$p((q, r) \rightarrow [-1]) := ([q + 1 + r] \rightarrow [0]).$$

In particular, p is given on objects by

$$p(q, r) := [q + 1 + r] \quad \text{and} \quad p(-1) := [0].$$

Definition 2.16 The (*generalized*) *path construction* is the induced functor

$$p^*: \mathcal{C}^{\Delta^{\text{op}}} \rightarrow \mathcal{C}^{\Sigma^{\text{op}}}.$$

Note that after composing with the underlying bisimplicial object functor i^* this path construction is the total décalage functor of [22].

Remark 2.17 Recall from Proposition 1.38 that for every $n \geq 0$ there is an isomorphism of augmented stable double categories

$$\mathcal{P}\Delta[n] \cong \mathcal{W}[n].$$

The analogous result in our new context is that there is an isomorphism of preaugmented bisimplicial sets

$$p^* \Delta[n] \cong \mathcal{W}[n],$$

where $\mathcal{W}[n]$ is as defined in Example 2.14. As usual, via Proposition 1.1 we can regard it as an isomorphism of discrete preaugmented bisimplicial objects in an arbitrary \mathcal{C} . This isomorphism can be verified directly from the definitions.

The functor p^* admits a right adjoint $p_*: \mathcal{C}^{\Sigma^{\text{op}}} \rightarrow \mathcal{C}^{\Delta^{\text{op}}}$ given by right Kan extension. Since, as we show later in Proposition 3.6, p^* generalizes the functor \mathcal{P} , we take the adjunction (\mathcal{P}, S_\bullet) as motivation for the following definition:

Definition 2.18 The (*generalized*) S_\bullet –*construction* is the right adjoint

$$p_*: \mathcal{C}^{\Sigma^{\text{op}}} \rightarrow \mathcal{C}^{\Delta^{\text{op}}}$$

to the generalized path construction.

Using Proposition 1.3, this adjoint pair has additional structure.

Proposition 2.19 The precomposition functor p^* and its right adjoint p_* form a \mathcal{C} –enriched adjunction

$$p^*: \mathcal{C}^{\Delta^{\text{op}}} \rightleftarrows \mathcal{C}^{\Sigma^{\text{op}}}: p_*.$$

Remark 2.20 The right adjoint

$$p_*: \mathcal{C}^{\Sigma^{\text{op}}} \rightarrow \mathcal{C}^{\Delta^{\text{op}}}$$

can be described explicitly. Indeed, by Proposition 1.3 and Remark 2.17, for every preaugmented bisimplicial object Y there are canonical isomorphisms

$$p_*(Y)_n \cong \text{Map}_{\mathcal{C}^{\Delta^{\text{op}}}}(\Delta[n], p_*Y) \cong \text{Map}_{\mathcal{C}^{\Sigma^{\text{op}}}}(p^* \Delta[n], Y) \cong \text{Map}_{\mathcal{C}^{\Sigma^{\text{op}}}}(\mathcal{W}[n], Y).$$

2.5 Augmented stable double Segal objects

In this section, we complete our definition of augmented stable double Segal objects. Since we have defined stable double Segal objects in the context of bisimplicial objects, we can use the underlying bisimplicial object functor i^* from Definition 2.10 to make sense of the double Segal condition and stability in the preaugmented setting.

Definition 2.21 A preaugmented bisimplicial object Y is *double Segal* if its underlying bisimplicial object i^*Y is double Segal. Similarly, it is *stable* if i^*Y is stable.

Mimicking the approach to 2–Segal objects from Proposition 1.17, we can express the double Segal property in terms of derived mapping spaces, using the representable preaugmented bisimplicial sets $\Sigma[q, r]$ from Example 2.12. This perspective will be useful when we develop our desired model structure. The following result is straightforward to check from the definitions:

Proposition 2.22 *The preaugmented bisimplicial object Y in \mathcal{C} is double Segal if and only if the maps*


$$\mathrm{Map}^h(\Sigma[q, r], Y) \rightarrow \mathrm{Map}^h(\Sigma[q, 1] \amalg_{\Sigma[q, 0]} \cdots \amalg_{\Sigma[q, 0]} \Sigma[q, 1], Y),$$
$$\mathrm{Map}^h(\Sigma[q, r], Y) \rightarrow \mathrm{Map}^h(\Sigma[1, r] \amalg_{\Sigma[0, r]} \cdots \amalg_{\Sigma[0, r]} \Sigma[1, r], Y)$$

induced by

$$\sigma^{q,r} : \Sigma[q, 1] \amalg_{\Sigma[q, 0]} \cdots \amalg_{\Sigma[q, 0]} \Sigma[q, 1] \rightarrow \Sigma[q, r],$$
$$\bar{\sigma}^{q,r} : \Sigma[1, r] \amalg_{\Sigma[0, r]} \cdots \amalg_{\Sigma[0, r]} \Sigma[1, r] \rightarrow \Sigma[q, r],$$

namely, the Segal maps in the second and first variable, respectively, are weak equivalences in \mathcal{C} .

We now define what it means for a preaugmented double Segal object to be augmented. To do so, we generalize notation from Example 2.13 as follows. For any $0 < k < n$, denote by $e^h : \Sigma[0, k] \rightarrow \Sigma[0, n]$ the identification of $\Sigma[0, k]$ as the k arrows at the end of the horizontal chain of n arrows present in $\Sigma[0, n]$, as highlighted in the following picture:



Similarly, we denote by $b^h: \Sigma[0, k] \rightarrow \Sigma[0, n]$ the identification of $\Sigma[0, k]$ as the k arrows at the *beginning* of the horizontal chain of n arrows present in $\Sigma[0, n]$. We define $e^v, b^v: \Sigma[k, 0] \rightarrow \Sigma[n, 0]$ similarly. We refer the reader to the discussion in [Example 2.12](#) in regard to the slight abuse of notation in treating these arrows as if they were in a category.

Definition 2.23 A preaugmented bisimplicial object Y is *augmented* if for any $q, r \geq 1$, the composites

$$(2.24) \quad Y_{q,0} \xrightarrow{t_v^h} Y_{0,0} \times Y_{-1} \xrightarrow{\text{pr}_1} Y_{q,0} \xrightarrow{b_v} Y_{q-1,0},$$

$$(2.25) \quad Y_{0,r} \xrightarrow{s_h^h} Y_{0,0} \times Y_{-1} \xrightarrow{\text{pr}_1} Y_{0,r} \xrightarrow{e_h} Y_{0,r-1}$$

are weak equivalences, where the maps b_v and e_h are induced by the maps of representables b^v and e^h .

We warn the reader that our notion of augmented bisimplicial object is different than the one used by Carlier² [10, Section 2.2].

Under the double Segal condition, to check whether a preaugmented bisimplicial object is augmented it suffices to consider the above maps only for $q = 1$ and $r = 1$.

Lemma 2.26 A preaugmented double Segal object Y is augmented if and only if the composites

$$(2.27) \quad Y_{1,0} \xrightarrow{t_v^h} Y_{0,0} \times Y_{-1} \xrightarrow{\text{pr}_1} Y_{1,0} \xrightarrow{b_v} Y_{0,0},$$

$$(2.28) \quad Y_{0,1} \xrightarrow{s_h^h} Y_{0,0} \times Y_{-1} \xrightarrow{\text{pr}_1} Y_{0,1} \xrightarrow{e_h} Y_{0,0}$$

are weak equivalences in \mathcal{C} .

Proof We show that having a weak equivalence in (2.27) implies having one in (2.24); a similar argument shows that having a weak equivalence in (2.28) implies having a weak equivalence in (2.25).

The composite map in (2.24),

$$Y_{q,0} \xrightarrow{t_v^h} Y_{0,0} \times Y_{-1} \xrightarrow{\text{pr}_1} Y_{q,0} \xrightarrow{b_v} Y_{q-1,0},$$

²There, an augmented bisimplicial ∞ -groupoid is an object in $\text{Fun}((\Delta_+ \times \Delta_+)^{\text{op}} \setminus \{(-1, -1)\}, \text{Spaces})$. Thus, they are similar to our preaugmented bisimplicial objects in ∞ -groupoids, but have many extra objects $X_{q,-1}$ and $X_{-1,r}$ rather than the single object X_{-1} , but without the extra condition we call the augmentation.

is the left vertical map in the homotopy commutative diagram (see [Remark 1.21](#))

$$\begin{array}{ccc} Y_{-1} \times_{Y_{0,0}}^{h_{t_v}} Y_{q,0} & \xrightarrow{\simeq} & Y_{-1} \times_{Y_{0,0}}^{h_{t_v}} \overbrace{(Y_{1,0} \times_{Y_{0,0}}^{s_v} \cdots \times_{Y_{0,0}}^{h_{t_v}} Y_{1,0})}^q \\ \downarrow & & \downarrow \simeq \\ & & (Y_{-1} \times_{Y_{0,0}}^{h_{t_v}} Y_{1,0}) \times_{Y_{0,0}}^{s_v} (Y_{1,0} \times_{Y_{0,0}}^{s_v} \cdots \times_{Y_{0,0}}^{h_{t_v}} Y_{1,0}) \\ & & \downarrow \simeq \\ Y_{q-1,0} & \xrightarrow{\simeq} & Y_{0,0} \times_{Y_{0,0}}^h \underbrace{(Y_{1,0} \times_{Y_{0,0}}^{s_v} \cdots \times_{Y_{0,0}}^{h_{t_v}} Y_{1,0})}_{q-1} \end{array}$$

The horizontal maps are given by Segal maps in the first index, and are equivalences since Y is double Segal. On the right, the top vertical map is given by associativity of the homotopy fiber product and the bottom one is the composition in [\(2.27\)](#). The two-out-of-three property finishes the proof. \square

As before, we can formulate the augmentation property in terms of derived mapping spaces, the proof of which we leave to the reader.

Proposition 2.29 *The preaugmented double Segal object Y is augmented if and only if the maps*

$$\begin{aligned} \mathrm{Map}^h(\Sigma[q, 0] \amalg_{\Sigma[0,0]} \Sigma[-1], Y) &\rightarrow \mathrm{Map}^h(\Sigma[q - 1, 0], Y), \\ \mathrm{Map}^h(\Sigma[0, r] \amalg_{\Sigma[0,0]} \Sigma[-1], Y) &\rightarrow \mathrm{Map}^h(\Sigma[0, r - 1], Y) \end{aligned}$$

induced by the inclusions

$$\begin{aligned} \bar{\zeta}^q : \Sigma[q - 1, 0] &\xrightarrow{b^v} \Sigma[q, 0] \rightarrow \Sigma[q, 0] \amalg_{\Sigma[0,0]} \Sigma[-1] = V[q], \\ \zeta^r : \Sigma[0, r - 1] &\xrightarrow{e^h} \Sigma[0, r] \rightarrow \Sigma[0, r] \amalg_{\Sigma[0,0]} \Sigma[-1] = H[r], \end{aligned}$$

respectively, are weak equivalences for all $q, r \geq 1$.

We give an analogous characterization for the stability condition.

Proposition 2.30 *A preaugmented double Segal object Y is stable if and only if the maps*

$$\begin{aligned} \mathrm{Map}^h(\Sigma[1, 1], Y) &\rightarrow \mathrm{Map}^h(\Sigma[1, 0] \amalg_{\Sigma[0,0]} \Sigma[0, 1], Y), \\ \mathrm{Map}^h(\Sigma[1, 1], Y) &\rightarrow \mathrm{Map}^h(\Sigma[0, 1] \amalg_{\Sigma[0,0]} \Sigma[1, 0], Y) \end{aligned}$$

$$\begin{aligned}\tau^{1,1}: \Sigma[1, 0] \amalg_{\Sigma[0,0]} \Sigma[0, 1] &\rightarrow \Sigma[1, 1], \\ \bar{\tau}^{1,1}: \Sigma[0, 1] \amalg_{\Sigma[0,0]} \Sigma[1, 0] &\rightarrow \Sigma[1, 1],\end{aligned}$$

In the former case, we include a cospan into a square, and in the latter case, we include a span into a square, as justified by the graphical explanations in (1.35).

Remark 2.31 Like augmented stable double categories, augmented stable double Segal objects enjoy nice symmetry properties. Specifically, if Y is an augmented stable double Segal object, then $Y_{1,0}$ and $Y_{0,1}$ are equivalent objects of \mathcal{C} . More precisely, there is a zigzag of weak equivalences connecting $Y_{1,0}$ and $Y_{0,1}$:

$$\begin{array}{c}
Y_{1,0} \simeq Y_{1,0} \overset{h}{\times}_{Y_{0,0}} Y_{0,0} \leftarrow^{\cong} Y_{1,0} \overset{h}{\times}_{Y_{0,0}} (Y_{0,1} \overset{h}{\times}_{Y_{0,0}} Y_{-1}) \leftarrow^{\cong} (Y_{1,0} \overset{h}{\times}_{Y_{0,0}} Y_{0,1}) \overset{h}{\times}_{Y_{0,0}} Y_{-1} \\
\phantom{Y_{1,0} \simeq Y_{1,0} \overset{h}{\times}_{Y_{0,0}} Y_{0,0} \leftarrow^{\cong} Y_{1,0} \overset{h}{\times}_{Y_{0,0}} (Y_{0,1} \overset{h}{\times}_{Y_{0,0}} Y_{-1}) \leftarrow^{\cong}} \quad \quad \quad \uparrow^{\cong} \\
\phantom{Y_{1,0} \simeq Y_{1,0} \overset{h}{\times}_{Y_{0,0}} Y_{0,0} \leftarrow^{\cong} Y_{1,0} \overset{h}{\times}_{Y_{0,0}} (Y_{0,1} \overset{h}{\times}_{Y_{0,0}} Y_{-1}) \leftarrow^{\cong}} \quad \quad \quad Y_{1,1} \overset{h}{\times}_{Y_{0,0}} Y_{-1} \\
\phantom{Y_{1,0} \simeq Y_{1,0} \overset{h}{\times}_{Y_{0,0}} Y_{0,0} \leftarrow^{\cong} Y_{1,0} \overset{h}{\times}_{Y_{0,0}} (Y_{0,1} \overset{h}{\times}_{Y_{0,0}} Y_{-1}) \leftarrow^{\cong}} \quad \quad \quad \downarrow^{\cong} \\
Y_{0,1} \simeq Y_{0,1} \overset{h}{\times}_{Y_{0,0}} Y_{0,0} \leftarrow^{\cong} Y_{0,1} \overset{h}{\times}_{Y_{0,0}} (Y_{1,0} \overset{h}{\times}_{Y_{0,0}} Y_{-1}) \leftarrow^{\cong} (Y_{0,1} \overset{h}{\times}_{Y_{0,0}} Y_{1,0}) \overset{h}{\times}_{Y_{0,0}} Y_{-1}
\end{array}$$

The horizontal maps are equivalences by the augmentation conditions (2.27) and (2.28) and by associativity of homotopy fiber products. The vertical maps are equivalences by stability.

With a similar argument, one can prove that for every $q \geq 0$ and $0 \leq i \leq q$, the spaces $Y_{q,0}$ and $Y_{q-i,i}$ are weakly equivalent.

3 Relationship with other S_\bullet -constructions

In this section, we exhibit two sources of examples for preaugmented bisimplicial object, quite different in nature. First, we define an augmented nerve for augmented stable double categories, enabling us to compare to the discrete version of the S_\bullet -construction developed in [6]. Second, we summarize a comparison to previously known S_\bullet -constructions; full details can be found in [7].

3.1 Comparison to augmented stable double categories

The examples of discrete augmented stable double Segal sets from the previous section all arise from augmented stable double categories via the following nerve construction, which is induced by the cosimplicial augmented stable double category $\mathcal{W}[\bullet]$.

Definition 3.1 The *augmented double nerve* of an augmented stable double category \mathcal{D} is the preaugmented bisimplicial set $N^a\mathcal{D}: \Sigma^{\text{op}} \rightarrow \text{Set}$ obtained by applying the functor $\text{Hom}_{\mathcal{D}\text{Cat}_{\text{aug}}^{\text{st}}}(-, \mathcal{D})$ to the composite

$$\Sigma \xrightarrow{p} \Delta \xrightarrow{\mathcal{W}[\bullet]} \mathcal{D}\text{Cat}_{\text{aug}}^{\text{st}}.$$

In other words, for $q, r \geq 0$,

$$N^a\mathcal{D}_{q,r} = \text{Hom}_{\mathcal{D}\text{Cat}_{\text{aug}}^{\text{st}}}(\mathcal{W}[p(q, r)], \mathcal{D}) = \text{Hom}_{\mathcal{D}\text{Cat}_{\text{aug}}^{\text{st}}}(\mathcal{W}[q + 1 + r], \mathcal{D}),$$

$$N^a\mathcal{D}_{-1} = \text{Hom}_{\mathcal{D}\text{Cat}_{\text{aug}}^{\text{st}}}(\mathcal{W}[p(-1)], \mathcal{D}) = \text{Hom}_{\mathcal{D}\text{Cat}_{\text{aug}}^{\text{st}}}(\mathcal{W}[0], \mathcal{D}) \cong \mathcal{A}\mathcal{D}.$$

The augmented nerve defines a functor

$$N^a: \mathcal{D}\text{Cat}_{\text{aug}}^{\text{st}} \rightarrow \text{Set}^{\Sigma^{\text{op}}}.$$

Let us consider the augmented nerves of our three families of examples of double categories.

Example 3.2 For any $n \geq 0$, the preaugmented bisimplicial space $\mathcal{W}[n]$ from [Example 2.14](#) is the augmented nerve of the double category $\mathcal{W}[n]$ from [Example 1.29](#), in the sense that there is an isomorphism

$$\mathcal{W}[n] \cong N^a(\mathcal{W}[n]).$$

The motivation for defining the augmented double nerve as such is that it agrees with the usual nerve of double categories from [\[14\]](#) when forgetting to the category $\mathcal{D}\text{Cat}$ of double categories, in the sense of the following proposition. Here, we denote the category of bisimplicial sets by $\text{Set}^{(\Delta \times \Delta)^{\text{op}}}$.

Proposition 3.3 The following diagram commutes up to natural isomorphism:

$$\begin{array}{ccc} \mathcal{D}\text{Cat}_{\text{aug}}^{\text{st}} & \xrightarrow{\text{forget}} & \mathcal{D}\text{Cat} \\ N^a \downarrow & & \downarrow N \\ \text{Set}^{\Sigma^{\text{op}}} & \xrightarrow{i^*} & \text{Set}^{(\Delta \times \Delta)^{\text{op}}} \end{array}$$


Proof Since the double nerve N is corepresented by the bi-cosimplicial double category $(q, r) \mapsto [q] \boxtimes [r]$ from [Example 1.34](#), it is enough to construct a bijection

$$(i^* N^a \mathcal{D})_{q,r} = \operatorname{Hom}_{\mathcal{D}Cat_{\text{aug}}^{\text{st}}}(\mathcal{W}[q+1+r], \mathcal{D}) \cong \operatorname{Hom}_{\mathcal{D}Cat}([q] \boxtimes [r], \mathcal{D}) = (N\mathcal{D})_{q,r}$$

that is natural in q and r . We construct a double functor

$$[q] \boxtimes [r] \rightarrow \mathcal{W}[q+1+r]$$

by sending the object (i, j) to $(i, j+q+1)$, and extending uniquely to horizontal morphisms, vertical morphisms and squares. For instance, the image of $[1] \boxtimes [2]$ into $\mathcal{W}[4]$ is highlighted in the following picture:



The functor induces the right-hand map in the diagram

$$\begin{array}{ccc} \operatorname{Hom}_{\mathcal{D}Cat_{\text{aug}}^{\text{st}}}(\mathcal{W}[q+1+r], \mathcal{D}) & & \operatorname{Hom}_{\mathcal{D}Cat}([q] \boxtimes [r], \mathcal{D}) \\ & \searrow & \nearrow \\ & \operatorname{Hom}_{\mathcal{D}Cat}(\mathcal{W}[q+1+r], \mathcal{D}) & \end{array}$$

Using stability, techniques similar to those used in [\[6, Proposition 4.9\]](#) can be used to show that the composite map is a bijection. Roughly speaking, given a diagram in \mathcal{D} indexed over the red portion of the picture, the image of 11 can be determined using the horizontal augmentation in \mathcal{D} , the image of 01 via stability, and the image of 00 again using the horizontal augmentation. The lower part of the diagram can be reconstructed in a similar way, but instead using the vertical direction of the augmentation and the opposite stability condition. □

The following proposition enables us to identify an augmented stable double category with the preaugmented bisimplicial set given by its augmented nerve:

Proposition 3.4 *The augmented nerve functor $N^a : \mathcal{DCat}_{\text{aug}}^{\text{st}} \rightarrow \text{Set}^{\Sigma^{\text{op}}}$ is fully faithful.*

Proof Given augmented stable double categories \mathcal{D} and \mathcal{E} , the commutative square

$$\begin{array}{ccc} \text{Hom}_{\mathcal{DCat}_{\text{aug}}^{\text{st}}}(\mathcal{D}, \mathcal{E}) & \xrightarrow{\text{forget}} & \text{Hom}_{\mathcal{DCat}}(\mathcal{D}, \mathcal{E}) \\ N^a \downarrow & & \downarrow N \\ \text{Hom}_{\text{Set}^{\Sigma^{\text{op}}}}(N^a \mathcal{D}, N^a \mathcal{E}) & \xrightarrow{i^*} & \text{Hom}_{\text{Set}^{(\Delta \times \Delta)^{\text{op}}}}(N \mathcal{D}, N \mathcal{E}) \end{array}$$

is a pullback square by definition of the augmented nerve. Since the nerve functor on double categories $N : \mathcal{DCat} \rightarrow \text{Set}^{(\Delta \times \Delta)^{\text{op}}}$ is fully faithful [13, Proposition 2.17], the right vertical arrow is an isomorphism, and therefore so is the left vertical arrow. Thus the augmented nerve $N^a : \mathcal{DCat}_{\text{aug}}^{\text{st}} \rightarrow \text{Set}^{\Sigma^{\text{op}}}$ is fully faithful, as desired. \square

We can now show that augmented stable double Segal sets indeed generalize augmented stable double categories via the augmented nerve construction.

Proposition 3.5 *If \mathcal{D} is an augmented stable double category, then its augmented nerve $N^a \mathcal{D}$ is an augmented stable double Segal set.*

Proof First, we show that the underlying bisimplicial set of $N^a \mathcal{D}$ is a double Segal set. By Proposition 3.3, the underlying bisimplicial set of $N^a \mathcal{D}$ is the usual double category nerve $N \mathcal{D}$, which is a double Segal set analogously to the fact that usual nerves of categories are Segal sets.

To show that $N^a \mathcal{D}$ is stable, it is straightforward to check the condition in Lemma 2.7 using (1.27). One can similarly verify that $N^a \mathcal{D}$ is augmented by verifying Lemma 2.26 using (1.28). \square

Via the augmented nerve, the equivalence from Theorem 1.37,

$$\mathcal{P} : \mathcal{U}2\text{Seg} \rightleftarrows \mathcal{DCat}_{\text{aug}}^{\text{st}} : S_{\bullet},$$

is compatible with the adjunction

$$p^* : \mathcal{C}^{\Delta^{\text{op}}} \rightleftarrows \mathcal{C}^{\Sigma^{\text{op}}} : p_*$$

in the sense of the following proposition:

Proposition 3.6 *The following diagrams commute up to natural isomorphism:*

$$\begin{array}{ccc} \mathcal{U}2Seg & \xrightarrow{\mathcal{P}} & \mathcal{D}Cat_{\text{aug}}^{\text{st}} \\ \downarrow & & \downarrow N^a \\ sSet & \xrightarrow{p^*} & Set^{\Sigma^{\text{op}}} \end{array} \quad \text{and} \quad \begin{array}{ccc} \mathcal{U}2Seg & \xleftarrow{S_{\bullet}} & \mathcal{D}Cat_{\text{aug}}^{\text{st}} \\ \downarrow & & \downarrow N^a \\ sSet & \xleftarrow{p_*} & Set^{\Sigma^{\text{op}}} \end{array}$$

In light of this compatibility, we henceforth use the following notation:

Notation 3.7 We denote by \mathcal{P} the path construction p^* from [Definition 2.16](#), and by S_{\bullet} the right adjoint p_* from [Definition 2.18](#).

Proof of Proposition 3.6 Let X be a 2-Segal set, and recall that it is necessarily unital. Commutativity of the first diagram follows from the bijections

$$\begin{aligned} (p^*X)_{q,r} &= X_{q+1+r} \\ &\cong \text{Hom}_{\mathcal{U}2Seg}(\Delta[q+1+r], X) \\ &\cong \text{Hom}_{\mathcal{D}Cat_{\text{aug}}^{\text{st}}}(\mathcal{P}(\Delta[q+1+r]), \mathcal{P}X) \\ &\cong \text{Hom}_{\mathcal{D}Cat_{\text{aug}}^{\text{st}}}(\mathcal{W}[q+1+r], \mathcal{P}X) \\ &= N^a(\mathcal{P}X)_{q,r}, \end{aligned}$$

which are justified as follows. The initial equality is the definition of p^* . The first isomorphism follows from corepresentability, while the second is given by fact that \mathcal{P} is fully faithful from [Theorem 1.37](#). The last isomorphism follows from [Proposition 1.38](#), and the last line is simply the definition of the augmented nerve. Similarly,

$$(p_*X)_{-1} = X_0 = A(\mathcal{P}X) = N^a(\mathcal{P}X)_{-1}.$$

Moreover, these bijections are natural in Σ .

In order to prove that the second diagram commutes, let \mathcal{D} be an augmented stable double category. There are bijections

$$\begin{aligned} (p_*N^a\mathcal{D})_n &\cong \text{Hom}_{sSet}(\Delta[n], p_*N^a\mathcal{D}) \\ &\cong \text{Hom}_{Set^{\Sigma^{\text{op}}}}(p^*\Delta[n], N^a\mathcal{D}) \\ &\cong \text{Hom}_{Set^{\Sigma^{\text{op}}}}(\mathcal{W}[n], N^a\mathcal{D}) \\ &\cong \text{Hom}_{Set^{\Sigma^{\text{op}}}}(N^a\mathcal{W}[n], N^a\mathcal{D}) \\ &\cong \text{Hom}_{\mathcal{D}Cat_{\text{aug}}^{\text{st}}}(\mathcal{W}[n], \mathcal{D}) \\ &= S_n(\mathcal{D}) \end{aligned}$$

which are natural in Δ . The first two isomorphisms arise from the Yoneda lemma and the adjunction (p^*, p_*) . The others are given by [Remark 2.17](#), [Example 3.2](#) and [Proposition 3.4](#), respectively. Thus we conclude that the diagram commutes up to isomorphism. \square

So far we have discussed how to view augmented stable double categories as augmented stable double Segal sets. However, one might ask whether the same is true for any choice of \mathcal{C} using the discrete functor $d_*: \mathcal{S}et^{\Sigma^{op}} \rightarrow \mathcal{C}^{\Sigma^{op}}$ from [Section 1.1](#). Note that, a priori, the image of d_* is merely a preaugmented bisimplicial object in \mathcal{C} . As the augmentation, double Segal and stability conditions are all defined in terms of homotopy pullbacks, a sufficient condition for preservation of these structures is that $d: \mathcal{S}et \rightarrow \mathcal{C}$ preserves homotopy pullbacks.³ It is not clear that this property holds for any \mathcal{C} , but we verify it for the case of $\mathcal{C} = s\mathcal{S}et$ in the following example, which is the most important example in practice.

Example 3.8 Since every map of discrete simplicial sets is a Kan fibration, the strict pullback of maps between discrete spaces yields a model for the homotopy pullback. Thus, for any diagram of discrete spaces

$$A \rightarrow C \leftarrow B,$$

the canonical map from the pullback to the homotopy pullback gives an equivalence

$$A \times_C B \xrightarrow{\simeq} A \mathbin{\mathop{\times}\limits^h_C} B.$$

In particular, this immediately implies the following comparison:


Proposition 3.9 *A preaugmented bisimplicial set is double Segal, augmented or stable if and only if it is such when regarded as a discrete preaugmented bisimplicial space.*

3.2 Preview of proto-exact $(\infty, 1)$ –categories

In this section we give an overview of how augmented stable double Segal spaces arise naturally from the inputs of previous S_\bullet –constructions, namely from proto-exact $(\infty, 1)$ –categories, which generalize both exact categories and stable $(\infty, 1)$ –categories. In particular, our generalized S_\bullet –construction is a natural generalization of ones which have been previously developed in these contexts. In the companion paper [\[7\]](#), we give detailed proofs of the statements mentioned here and in particular look closely at the special cases of exact categories and stable $(\infty, 1)$ –categories.

³The appendix of [\[18\]](#) gives sufficient conditions for d to preserve finite limits and hence pullbacks.

We summarize these different settings and constructions in the following diagram:



Roughly speaking, we start with a homotopical category, together with extra data which distinguishes certain objects, morphisms and commutative squares. An appropriate nerve functor N^{pe} assigns distinguished objects to the object $[-1]$ of Σ and arrays of size $q \times r$ of commuting squares to the object (q, r) .

While we do not want to go into the details of $(\infty, 1)$ –category theory here, the main idea is that an $(\infty, 1)$ –category should be thought of as a category up to homotopy, with a discrete collection of objects but with mapping spaces for which composition is defined only up to homotopy. While there are several different ways to model such a structure, here we use the framework of quasicategories, which has been developed by Joyal [23] and Lurie [27]. Quasicategories are simplicial sets which satisfy certain horn-filling conditions. They are more general than Kan complexes, which can be thought of as groupoids up to homotopy. Often in the literature, the term “ ∞ –category” is used rather than “quasicategory”, but we use the latter for the sake of precision.

Let us now turn to the notion of proto-exact quasicategory, which was defined by Dyckerhoff and Kapranov [11, Definition 7.2.1]. Although they simply called them “exact”, we use this name to distinguish them from the related, but not identical, exact quasicategories in the sense of Barwick [4; 5], which are special cases of ours. The idea is that this structure includes that of an exact category as a special case, yet is general enough to include interesting examples that do not fit into the previous framework, such as sub-quasicategories of a stable quasicategory which are closed under extensions, as considered in [5]. The terminology used is very similar to what is used in the definition of an exact category, so a reader only familiar with that level of generality can work in that context instead.

Definition 3.10 A *proto-exact quasicategory* consists of a triple of quasicategories $(\mathcal{Q}, \mathcal{M}, \mathcal{E})$ such that

- both \mathcal{M} and \mathcal{E} are sub-quasicategories of \mathcal{Q} containing all equivalences;

- the category \mathcal{Q} has a zero object, \mathcal{M} contains all the morphisms whose source is a zero object, and \mathcal{E} contains all the morphisms whose target is a zero object;
- any pushout of a morphism in \mathcal{M} along a morphism of \mathcal{E} exists and belongs to \mathcal{M} and any pushout of a morphism in \mathcal{E} along a morphism of \mathcal{M} is in \mathcal{E} , and similarly for pullbacks; and
- a square whose horizontal morphisms are in \mathcal{M} and whose vertical morphisms are in \mathcal{E} is cartesian if and only if it is cocartesian.

A morphism between proto-exact quasicategories preserves the appropriate structure. We now define a nerve functor which takes a proto-exact quasicategory to a preaugmented bisimplicial space. For any quasicategory \mathcal{Q} , we denote by $J(\mathcal{Q})$ the maximal Kan complex spanned by its vertices, as in [24, Proposition 1.16], and by $\mathcal{Q}^{[q] \times [r]}$ the quasicategory of functors $[q] \times [r] \rightarrow \mathcal{Q}$.

Definition 3.11 The *proto-exact nerve* $N^{\text{pe}}\mathcal{Q}$ of a proto-exact quasicategory $\mathcal{Q} = (\mathcal{Q}, \mathcal{M}, \mathcal{E})$ is the preaugmented bisimplicial space $N^{\text{pe}}\mathcal{Q}: \Sigma^{\text{op}} \rightarrow s\text{Set}$ defined as follows:

- (1) The component in degree (q, r) is the simplicial set

$$N^{\text{pe}}\mathcal{Q}_{q,r} \subset J(\mathcal{Q}^{[q] \times [r]})$$

spanned by $q \times r$ grids in \mathcal{Q} with horizontal morphisms in \mathcal{M} , vertical morphisms in \mathcal{E} and all squares bicartesian.

- (2) The augmentation space is the simplicial set

$$N^{\text{pe}}\mathcal{Q}_{-1} \subset J(\mathcal{Q})$$

spanned by all the zero objects of \mathcal{Q} .

The bisimplicial structure is induced by the bi-cosimplicial structure of the categories $[q] \times [r]$. The additional map is the canonical inclusion of the space of zero objects into \mathcal{Q} .

As we prove in [7], we can say more.

Proposition 3.12 The proto-exact nerve N^{pe} defines a functor from the category of proto-exact quasicategories to the category of augmented stable double Segal spaces.

The S_{\bullet} -construction for (proto)-exact quasicategories was considered by Barwick [4; 5] and by Dyckerhoff and Kapranov [11, Section 2.4], and we denote the result of applying

this construction to a proto-exact quasicategory \mathcal{Q} by $S_{\bullet}(\mathcal{Q})$. The main result of [7] is the following theorem:

Theorem 3.13 *Let \mathcal{Q} be a proto-exact quasicategory. There is a levelwise weak equivalence of 2–Segal spaces*

$$S_{\bullet}(\mathcal{Q}) \xrightarrow{\cong} S_{\bullet}(N^{\text{pc}}\mathcal{Q}).$$

4 The model structures

Now that we have introduced the categories of objects we would like to work with, we equip them with homotopical structure. Specifically, we define model structures for 2–Segal objects and for augmented stable double Segal objects which encode the homotopy theories we are interested in.

4.1 The model structure for 2–Segal spaces

Following the approach of Dyckerhoff and Kapranov [11, Section 5.3] we construct a model structure for 2–Segal spaces via an application of Theorem 1.11. Here we use the mapping space description of 2–Segal spaces from Proposition 1.17 to determine which maps to use for the localization.

Definition 4.1 Let \mathcal{S} be the set consisting of the maps

$$f_P: \Delta[P] \rightarrow \Delta[n],$$

where P is a triangulation of the $(n+1)$ –gon and $\Delta[P]$ is as in Notation 1.16.

We now localize the injective model structure on $\mathcal{C}^{\Delta^{\text{op}}}$ with respect to this set \mathcal{S} , using Theorem 1.11.

Proposition 4.2 [11, Section 5.3] *Localizing the injective model structure on $\mathcal{C}^{\Delta^{\text{op}}}$ with respect to the set \mathcal{S} results in a model structure, which we denote by $\mathcal{C}_{\mathcal{S}}^{\Delta^{\text{op}}}$, in which the fibrant objects are precisely the injectively fibrant 2–Segal objects.*

Observe that, by Theorem 1.25, the fibrant objects of this model structure satisfy unitality. The following result, which is used in the proof of Lemma 6.4, provides a convenient reformulation of the fact that every 2–Segal object is unital. Specifically, the map $\omega_{n,i}$ described below is one of the ways to encode unitality, and we establish that is a weak equivalence in the localized model structure $\mathcal{C}_{\mathcal{S}}^{\Delta^{\text{op}}}$.

Proposition 4.3 *The map*

$$\omega_{n,i}: \Delta[n-1] \xrightarrow{d^i} \Delta[n] \rightarrow \Delta[n] \amalg_{\Delta[1]}^{\beta^i, s^0} \Delta[0],$$

where $n \geq 0$, $0 \leq i \leq n$ and the map β^i is as in [Definition 1.22](#), is a weak equivalence in the localized model structure $\mathcal{C}_S^{\Delta^{\text{op}}}$.

Proof We first observe that the map $\omega_{n,i}$ is right inverse to the map [\(1.24\)](#),

$$\Delta[n] \amalg_{\Delta[1]}^{\beta^i, s^0} \Delta[0] \rightarrow \Delta[n-1],$$

from [Remark 1.23](#). Therefore, it is enough to show that the latter map is a weak equivalence. To this end, let X be an injectively fibrant 2-Segal object, and we look at the induced map

$$X_{n-1} \simeq \text{Map}^h(\Delta[n-1], X) \rightarrow \text{Map}^h(\Delta[n] \amalg_{\Delta[1]}^{\beta^i, s^0} \Delta[0], X) \simeq X_n \times_{X_1}^h X_0.$$

We know by [Remark 1.23](#) that this map is a weak equivalence when X is unital, which we know by [Theorem 1.25](#). □

4.2 The model structure for augmented stable double Segal objects

The model structure for augmented stable double Segal objects is also obtained via localization. The maps with respect to which we localize are those developed in [Propositions 2.22](#), [2.29](#) and [2.30](#), and those results demonstrate that the local objects have precisely the desired properties.

Definition 4.4 Let \mathcal{T} be the union of the following three sets of maps in $\mathcal{C}^{\Sigma^{\text{op}}}$:

- (1) the set $\mathcal{T}_{\text{Segal}}$ of maps

$$\sigma^{q,r}: \Sigma[q, 1] \amalg_{\Sigma[q,0]} \cdots \amalg_{\Sigma[q,0]} \Sigma[q, 1] \rightarrow \Sigma[q, r],$$

induced by the Segal map in the second variable, and

$$\bar{\sigma}^{q,r}: \Sigma[1, r] \amalg_{\Sigma[0,r]} \cdots \amalg_{\Sigma[0,r]} \Sigma[1, r] \rightarrow \Sigma[q, r],$$

induced by the Segal map in the first variable, each for all $q, r \geq 2$;

- (2) the set \mathcal{T}_{aug} of maps

$$\zeta^r: \Sigma[0, r-1] \rightarrow \Sigma[0, r] \rightarrow \Sigma[0, r] \amalg_{\Sigma[0,0]} \Sigma[-1] = H[r]$$

for every r , induced by the map b^h , and

$$\bar{\zeta}^q: \Sigma[q-1, 0] \rightarrow \Sigma[q, 0] \rightarrow \Sigma[q, 0] \amalg_{\Sigma[0,0]} \Sigma[-1] = V[q]$$

for every $q \geq 0$, induced by the map e^v ; and

(3) the set $\mathcal{T}_{\text{stable}}$ of maps

$$\tau^{q,r} : \Sigma[q, 0] \amalg_{\Sigma[0,0]} \Sigma[0, r] \rightarrow \Sigma[q, r],$$

induced by the inclusion of the cospan, and

$$\bar{\tau}^{q,r} : \Sigma[0, r] \amalg_{\Sigma[0,0]} \Sigma[q, 0] \rightarrow \Sigma[q, r],$$

induced by the inclusion of the span, each for all $q, r \geq 0$.

Similarly to before, we can localize the injective model structure on $\mathcal{C}^{\Sigma^{\text{op}}}$ with respect to the set \mathcal{T} using [Theorem 1.11](#).

Proposition 4.5 *Localizing the injective model structure on $\mathcal{C}^{\Sigma^{\text{op}}}$ results in a model structure, which we denote by $\mathcal{C}_{\mathcal{T}}^{\Sigma^{\text{op}}}$, in which the fibrant objects are precisely the augmented stable double Segal objects which are injectively fibrant.*

4.3 The auxiliary model structure on $\mathcal{C}^{\Sigma^{\text{op}}}$

Our goal is to prove that the model categories $\mathcal{C}_{\mathcal{S}}^{\Delta^{\text{op}}}$ and $\mathcal{C}_{\mathcal{T}}^{\Sigma^{\text{op}}}$ are Quillen equivalent. However, for the proof we use an intermediate model structure on $\mathcal{C}^{\Sigma^{\text{op}}}$ which is easier to compare to $\mathcal{C}_{\mathcal{S}}^{\Delta^{\text{op}}}$. Once again, this model structure is given by a localization of the injective model structure on $\mathcal{C}^{\Sigma^{\text{op}}}$.

The idea behind this localization is to apply the path construction $\mathcal{P} = p^*$ to the maps in \mathcal{S} . With the isomorphism from [Proposition 1.38](#) in mind, we make the following definition:


Definition 4.6 Let P be a polygonal decomposition of the regular $(n+1)$ –gon, and recall the simplicial set $\Delta[P]$ from [Notation 1.16](#). Let $\mathcal{W}[P]$ be the preaugmented bisimplicial set

$$\mathcal{W}[P] := \mathcal{P}\Delta[P].$$

In particular, when P is a decomposition of the $(n+1)$ –gon into a triangle and an n –gon, $\mathcal{W}[P]$ can be expressed as a pushout of objects

$$\mathcal{W}[P] = \mathcal{P}(\Delta[P]) = \mathcal{P}(\Delta[2] \amalg_{\Delta[1]} \Delta[n-1]) \cong \mathcal{W}[2] \amalg_{\mathcal{W}[1]} \mathcal{W}[n-1].$$

Below is an image of $W[P]$ in the case where $n = 4$ and P is the decomposition of a pentagon into a triangle and a quadrilateral by adding the diagonal from 0 to 2:



Treating $W[P]$ as a discrete object in $\mathcal{C}^{\Sigma^{\text{op}}}$ for any \mathcal{C} , we can now define the maps with respect to which we want to localize.

Definition 4.7 Let \mathcal{W} be the set of maps, in $\mathcal{C}^{\Sigma^{\text{op}}}$,

$$\mathcal{P}f_P: \mathcal{P}\Delta[P] \cong W[P] \rightarrow W[n] \cong \mathcal{P}\Delta[n],$$

where P is a triangulation of the $(n+1)$ -gon.

As in the previous two localizations, we apply [Theorem 1.11](#) to obtain the following model structure:

Proposition 4.8 *Localizing the injective model structure on $\mathcal{C}^{\Sigma^{\text{op}}}$ with respect to \mathcal{W} results in a model structure which we denote by $\mathcal{C}_{\mathcal{W}}^{\Sigma^{\text{op}}}$.*

Remark 4.9 In the next section, we show that every \mathcal{T} -local object in $\mathcal{C}^{\Sigma^{\text{op}}}$ is also \mathcal{W} -local, a key fact in comparing the two model structures on $\mathcal{C}^{\Sigma^{\text{op}}}$. However, the converse result fails in general, showing that the two localizations do not coincide. Specifically, for $\mathcal{C} = sSet$, we claim that $\Sigma[0, 0]$ is \mathcal{W} -local but not \mathcal{T} -local in $sSet^{\Sigma^{\text{op}}}$. To see that $\Sigma[0, 0]$ is \mathcal{W} -local, first observe that, for any n and any decomposition P , we have that

$$\text{Map}(W[n], \Sigma[0, 0]) = \text{Map}(W[P], \Sigma[0, 0]) = \varnothing,$$

because the augmentation of $\Sigma[0, 0]$ is empty, whereas the augmentations of $W[n]$ and $W[P]$ are not. However, one can check that $\Sigma[0, 0]$ is injectively fibrant, so the isomorphism between the nonderived mapping spaces induces a weak equivalence of derived mapping spaces. As a consequence $\Sigma[0, 0]$ is \mathcal{W} -local.

To see that $\Sigma[0, 0]$ is not \mathcal{T} –local, consider the map

$$\zeta^1: \Sigma[0, 0] \hookrightarrow H[1]$$

in \mathcal{T} . Mapping into $\Sigma[0, 0]$, we get the map

$$\emptyset = \mathrm{Map}^h(H[1], \Sigma[0, 0]) \rightarrow \mathrm{Map}^h(\Sigma[0, 0], \Sigma[0, 0]) \neq \emptyset,$$

which cannot be induced by a weak equivalence.

5 The generalized S_\bullet –construction as a right Quillen functor

In this section we prove the following result, which is at the core of the comparison of our model structures. Recall the functors \mathcal{P} and S_\bullet from [Notation 3.7](#).

Theorem 5.1 *The path construction and the S_\bullet –construction induce a Quillen pair*

$$\mathcal{P}: \mathcal{C}_S^{\Delta^{\mathrm{op}}} \rightleftarrows \mathcal{C}_{\mathcal{T}}^{\Sigma^{\mathrm{op}}} : S_\bullet.$$

To prove this theorem, we make use of the auxiliary model structure, introduced in [Section 4.3](#), and, more specifically, we show that the two adjoint pairs

$$\mathcal{C}_S^{\Delta^{\mathrm{op}}} \xrightleftharpoons[S_\bullet]{\mathcal{P}} \mathcal{C}_{\mathcal{W}}^{\Sigma^{\mathrm{op}}} \xrightleftharpoons[\mathrm{id}]{\mathrm{id}} \mathcal{C}_{\mathcal{T}}^{\Sigma^{\mathrm{op}}}$$

are Quillen pairs, with the left adjoints written topmost.

We first establish the first of these Quillen adjunctions, since it is the easier of the two.

Proposition 5.2 *The path construction induces a Quillen pair*

$$\mathcal{P}: \mathcal{C}^{\Delta^{\mathrm{op}}} \rightleftarrows \mathcal{C}^{\Sigma^{\mathrm{op}}} : S_\bullet,$$

on injective model structures, which descends to a Quillen pair after localization,

$$\mathcal{P}: \mathcal{C}_S^{\Delta^{\mathrm{op}}} \rightleftarrows \mathcal{C}_{\mathcal{W}}^{\Sigma^{\mathrm{op}}} : S_\bullet.$$

Proof Let X be a simplicial object in \mathcal{C} . Using the definition of $\mathcal{P} = p^*$, recall that $(\mathcal{P}X)_{q,r} = X_{q+r+1}$ and $(\mathcal{P}X)_{-1} = X_0$. It follows that \mathcal{P} preserves levelwise cofibrations and levelwise weak equivalences and therefore is a left Quillen functor for the injective model structures by [\[20, Proposition 8.5.3\]](#).

To show that this Quillen pair is compatible with the localizations, recall that we defined \mathcal{W} precisely to be $\mathcal{P}(\mathcal{S})$. It follows that localizing $\mathcal{C}^{\Delta^{\mathrm{op}}}$ with respect to \mathcal{S} , and $\mathcal{C}^{\Sigma^{\mathrm{op}}}$ with respect to $\mathcal{P}(\mathcal{S})$, retains the structure of the Quillen pair by [\[20, Theorem 3.3.20\]](#). \square

Thus, we have reduced [Theorem 5.1](#) to comparing the two different model structures on $\mathcal{C}^{\Sigma^{\text{op}}}$, as we restate here.

Proposition 5.3 *The identity functors induce a Quillen pair*

$$\mathcal{C}_{\mathcal{W}}^{\Sigma^{\text{op}}} \rightleftarrows \mathcal{C}_{\mathcal{T}}^{\Sigma^{\text{op}}}.$$

Since its proof is quite involved, for organizational purposes we first give the main ingredients in [Section 5.1](#) and then in [Section 5.2](#) we return to some deferred technical points.

5.1 Main outline of the proof

Since cofibrations are the same in both model structures, the left adjoint, as the identity functor, preserves them. Thus, our strategy to complete the proof is to verify that every \mathcal{W} –local equivalence is a \mathcal{T} –local equivalence.

It suffices to show that the right adjoint, also the identity, preserves fibrant objects. Indeed, if A is \mathcal{T} –local, and hence \mathcal{W} –local, and $f: Y \rightarrow Y'$ is a \mathcal{W} –local equivalence, it follows that the induced map

$$\text{Map}^h(Y', A) \rightarrow \text{Map}^h(Y, A)$$

is a weak equivalence in \mathcal{C} , and therefore f is also a \mathcal{T} –local equivalence. Thus, to complete the proof of [Theorem 5.1](#) we only need the following result, whose proof is technical and occupies the remainder of the section:

Proposition 5.4 *Any \mathcal{T} –local object of $\mathcal{C}^{\Sigma^{\text{op}}}$ is \mathcal{W} –local.*

The key step is to prove that S_{\bullet} preserves fibrant objects, as made precise by the following proposition:

Proposition 5.5 *If Y is a \mathcal{T} –local object in $\mathcal{C}^{\Sigma^{\text{op}}}$, then $S_{\bullet}Y$ is \mathcal{S} –local.*

We already know from [Proposition 5.2](#) that S_{\bullet} preserves injectively fibrant objects. Therefore, to prove [Proposition 5.5](#), it suffices to show that $S_{\bullet}(Y)$ is a 2–Segal object and therefore \mathcal{S} –local.

To prove this result, which is the content of the next proposition, we need to know that certain maps are acyclic cofibrations in $\mathcal{C}_{\mathcal{T}}^{\Sigma^{\text{op}}}$. To improve readability we have deferred these results to [Section 5.2](#).

Proposition 5.6 *If an object Y of $\mathcal{C}^{\Sigma^{\text{op}}}$ is \mathcal{T} –local, then $S_{\bullet}Y$ is a 2–Segal object.*

Proof This argument is inspired by the proof of [11, Proposition 2.4.8]. Recall from Proposition 1.18 that to check 2–Segality it suffices to consider a decomposition P which is either the decomposition of the $(n+1)$ –gon into a triangle $\{0, 1, 2\}$ and the remaining n –gon, or the decomposition into a triangle $\{n-2, n-1, n\}$ and the remaining n –gon.

Consider the former case; the other decomposition can be treated similarly. We need to check that the 2–Segal map

$$\text{Map}^h(\Delta[n], S_{\bullet}Y) \rightarrow \text{Map}^h(\Delta[P], S_{\bullet}Y),$$

with the derived mapping spaces taken in the injective model structure, is a weak equivalence.

Let us begin by looking at similar derived mapping spaces for the original object Y . We prove in Lemma 5.9 that the inclusion $W[P] \hookrightarrow W[n]$ is an acyclic cofibration, and in particular a weak equivalence, in $\mathcal{C}_{\mathcal{T}}^{\Sigma^{\text{op}}}$. Since Y is \mathcal{T} –local, by definition the induced map

$$\text{Map}^h(W[n], Y) \rightarrow \text{Map}^h(W[P], Y)$$

is a weak equivalence. Moreover, since Y is injectively fibrant and all objects are cofibrant, this weak equivalence can be realized by underived mapping spaces

$$\text{Map}_{\mathcal{C}^{\Sigma^{\text{op}}}}(W[n], Y) \rightarrow \text{Map}_{\mathcal{C}^{\Sigma^{\text{op}}}}(W[P], Y).$$

Applying the isomorphisms $\mathcal{P}\Delta[n] \cong W[n]$ of Remark 2.17 and the definition of $W[P]$, we can rewrite this weak equivalence as

$$\text{Map}_{\mathcal{C}^{\Sigma^{\text{op}}}}(\mathcal{P}\Delta[n], Y) \rightarrow \text{Map}_{\mathcal{C}^{\Sigma^{\text{op}}}}(\mathcal{P}\Delta[P], Y),$$

to which we can apply the adjunction $(\mathcal{P}, S_{\bullet})$ to obtain the weak equivalence

$$\text{Map}_{\mathcal{C}^{\Delta^{\text{op}}}}(\Delta[n], S_{\bullet}Y) \rightarrow \text{Map}_{\mathcal{C}^{\Delta^{\text{op}}}}(\Delta[P], S_{\bullet}Y).$$

By inspection, this map is precisely the one induced by $f_P: \Delta[P] \rightarrow \Delta[n]$ from Definition 4.1. Thus, it only remains to verify that $S_{\bullet}(Y)$ is injectively fibrant, so that this map realizes the corresponding map on derived mapping spaces, which is precisely the 2–Segal map. However, Y is injectively fibrant as a \mathcal{T} –local object, and therefore $S_{\bullet}(Y)$ is also injectively fibrant by Proposition 5.2. We conclude that $S_{\bullet}Y$ is 2–Segal. \square

Now that we have proved [Proposition 5.5](#), we use it to prove [Proposition 5.4](#).

Proof of Proposition 5.4 Let Y be a \mathcal{T} –local object and let P be the decomposition of the $(n+1)$ –gon into a triangle $\{0, 1, 2\}$ and the remaining n –gon or into a triangle $\{n-2, n-1, n\}$ and the remaining n –gon, as in [Proposition 1.18](#). Consider the map

$$f_P \colon \Delta[P] \rightarrow \Delta[n]$$

corresponding to this decomposition. We want to prove that Y is $\{\mathcal{P}f_P\}$ –local.

Since $S_\bullet Y$ is \mathcal{S} –local by [Proposition 5.5](#), the map

$$(f_P)^* \colon \mathrm{Map}^h(\Delta[n], S_\bullet Y) \rightarrow \mathrm{Map}^h(\Delta[P], S_\bullet Y)$$

is a weak equivalence. Using similar manipulations between derived and underived mapping spaces and the adjunction (\mathcal{P}, S_\bullet) as in the proof of [Proposition 5.6](#), we obtain that the map


$$(\mathcal{P}f_P)^* \colon \mathrm{Map}^h(W[n], Y) \rightarrow \mathrm{Map}^h(W[P], Y)$$

is a weak equivalence, as desired. □

5.2 Acyclicity results

In this section, we return to some technical results that were deferred, most importantly [Lemma 5.9](#), which was used in the proof of [Proposition 5.6](#).

Lemma 5.7 *For $i = 0, 1$, the map $f_i \colon W[1] \xrightarrow{\mathcal{P}d^1} W[2] \rightarrow W[2] \amalg_{W[1]}^i W[0]$, whose image is displayed with red dotted arrows in*




for $i = 0, 1$, respectively, is an acyclic cofibration in $\mathcal{C}_{\mathcal{T}}^{\Sigma^{\mathrm{op}}}$.

Proof We prove that f_1 is an acyclic cofibration; the proof for f_0 is similar. Write f_1 as the composite

$$W[1] \hookrightarrow H[1] \amalg_{\Sigma[0,0]} W[2] \amalg_{W[1]}^1 W[0] \rightarrow W[2] \amalg_{W[1]}^1 W[0],$$


depicted by




We claim that both are weak equivalences.

The first map is obtained by the following several steps: First we fill the cospan formed by the unique vertical morphism and the identity on the target to a square. Then we add a horizontal augmentation for the newly added object. Finally, we add a horizontal composite of the augmentation map and the top vertical source in the square. Each of these steps is implemented by taking a pushout along an acyclic cofibration, and the resulting inclusion is therefore a \mathcal{T} –local equivalence.

The second map is obtained by identifying the copy of $H[1]$ at the top with the composition of the two horizontal maps in the copy of $W[2]$. It can be depicted as



It is a left inverse of the inclusion $H[1] \hookrightarrow H[1] \amalg_{\Sigma[0,0]} H[1]$ depicted by



which is a \mathcal{T} –local equivalence. Applying the two-out-of-three property finishes the proof. \square

For the remainder of this section, we prove acyclicity results for maps induced by decomposing polygons. Let us first set some notation.

Notation 5.8 Let P be the decomposition of the $(n+1)$ –gon into the triangle $\{0, 1, 2\}$ and the remaining n –gon, as in [Proposition 1.18](#). Denote by $H[P]$ the first row of $W[P]$, namely the preaugmented bisimplicial set given by

$$H[P] := W[P] \cap H[n] \subset W[n].$$

The description of $W[P]$ as a pushout from [Definition 4.6](#) restricts to an isomorphism

$$H[P] \cong H[2] \amalg_{H[1]} H[n-1].$$

Lemma 5.9 For any decomposition P as described above, the inclusion

$$W[P] \hookrightarrow W[n]$$

is an acyclic cofibration in $\mathcal{C}_{\mathcal{T}}^{\Sigma^{\text{op}}}$.

The proof of [Lemma 5.9](#) boils down to the following two technical results and their respective duals, whose proofs are similar.

Lemma 5.10 *Let P be the decomposition of the $(n+1)$ -gon into a triangle $\{0, 1, 2\}$ and the remaining n -gon. The inclusion $H[P] \hookrightarrow H[n]$ is an acyclic cofibration in $\mathcal{C}_{\mathcal{T}}^{\Sigma^{\text{op}}}$.*

Proof We want to prove that the canonical inclusion $H[P] \hookrightarrow H[n]$ is an acyclic cofibration by realizing it as a pushout along an acyclic cofibration. First, a standard argument for Segal objects can be used to show that the generalized Segal maps

$$(b^h, e^h): \Sigma[0, k] \amalg_{\Sigma[0, 0]} \Sigma[0, n - k] \rightarrow \Sigma[0, n]$$

are \mathcal{T} -local equivalences.

Then we observe that $\Sigma[0, n]$ contains a copy of $\Sigma[0, n - 1]$, highlighted in red, and a copy of $\Sigma[0, 2]$ given highlighted in dashed blue, which overlap on a copy of $\Sigma[0, 1]$,



The induced morphism

$$\Sigma[0, 2] \amalg_{\Sigma[0, 1]} \Sigma[0, n - 1] \rightarrow \Sigma[0, n]$$

is an acyclic cofibration, as can be observed from the commutative diagram

$$\begin{array}{ccc} \Sigma[0, 2] \amalg_{\Sigma[0, 1]} \Sigma[0, n - 1] & \xrightarrow{\quad\quad\quad} & \Sigma[0, n] \\ \uparrow \int \text{id} \amalg (b^h, e^h) & & \uparrow \int (b^h, e^h) \simeq \\ \Sigma[0, 2] \amalg_{\Sigma[0, 1]} \Sigma[0, 1] \amalg_{\Sigma[0, 0]} \Sigma[0, n - 2] & \xrightarrow{\cong} & \Sigma[0, 2] \amalg_{\Sigma[0, 0]} \Sigma[0, n - 2] \end{array}$$

where the vertical morphisms are induced by generalized Segal maps in the second variable and the bottom map is a canonical isomorphism.

Recall that $H[P] \cong H[2] \amalg_{H[1]} H[n - 1]$. Since there is an inclusion $\Sigma[0, k] \hookrightarrow H[k] = \Sigma[0, k] \amalg_{\Sigma[0, 0]} H[0]$ for every k , these maps induce the top horizontal map in the following pushout square:

$$\begin{array}{ccc} \Sigma[0, 2] \amalg_{\Sigma[0, 1]} \Sigma[0, n - 1] & \xrightarrow{\quad\quad\quad} & H[2] \amalg_{H[1]} H[n - 1] \\ \downarrow & & \downarrow \\ \Sigma[0, n] & \xrightarrow{\quad\quad\quad} & H[n] \end{array}$$

Note that both horizontal maps are the identity on the underlying bisimplicial subsets of $\Sigma[0, n]$. In particular, the canonical inclusion $H[P] \hookrightarrow H[n]$ is a pushout along an acyclic cofibration and hence is an acyclic cofibration itself. \square

The second fact we need is an analogue of [11, Lemma 2.4.9].


Lemma 5.11 *The inclusion $H[n] \hookrightarrow W[n]$ is an acyclic cofibration $\mathcal{C}_J^{\Sigma^{\text{op}}}$.*

Proof We prove by induction on $n \geq 0$ that $W[n]$ can be built out of $H[n]$ by taking pushouts along basic acyclic cofibrations. When $n = 0$, the inclusion $H[0] \hookrightarrow W[0]$ is the identity of $\Sigma[-1]$, so there is nothing to prove.

Now suppose $n > 0$. Define a filtration of preaugmented bisimplicial sets

$$H[n] \subseteq F^{(0)} \subseteq F^{(1)} \subseteq \dots \subseteq F^{(n)} \cong W[n]$$

in which all the inclusions are acyclic cofibrations. For $n = 4$, this filtration can be depicted as follows:



Let us describe this filtration more precisely:

- Define $F^{(0)}$ to be the preaugmented bisimplicial set obtained from $H[n]$ by adjoining a copy of $W[n-1]$ along the first $n-1$ arrows. More precisely, $F^{(0)}$ can be written as a pushout

$$F^{(0)} := W[n-1] \amalg_{H[n-1]} H[n],$$

which is depicted as in the codomain of the map j_0 above when $n = 4$. The inclusion of $H[n]$ into $F^{(0)}$ can be written in the form

$$H[n] \cong H[n-1] \amalg_{H[n-1]} H[n] \hookrightarrow W[n-1] \amalg_{H[n-1]} H[n] = F^{(0)},$$

which is an acyclic cofibration by our inductive hypothesis.

- For each $1 \leq m \leq n-1$, we obtain $F^{(m)}$ from $F^{(m-1)}$ by adjoining a new square and all horizontal and vertical composites with preexisting squares. When $n = 4$, this process can be visualized via the maps j_1 , j_2 and j_3 in the diagram above.

In order to build $F^{(m)}$ explicitly, we need several steps: (1) complete a span to a square, (2) add its horizontal composites, (3) add its vertical composites, and (4) make sure that the interchange law holds. We use a further filtration

$$F^{(m-1)} \subseteq F^{(m-1,1)} \subseteq F^{(m-1,2)} \subseteq F^{(m-1,3)} \subseteq F^{(m-1,4)} =: F^{(m)},$$

to encode these steps, in such a way that the inclusions are all acyclic cofibrations. We now describe this filtration more precisely:

- (1) To obtain $F^{(m-1,1)}$ from $F^{(m-1)}$, we complete the relevant span to a square, obtaining the m^{th} square in the last column. (We refer back to the above diagrams to identify the appropriate spans to the right.) More precisely, it is given as a pushout

$$F^{(m-1,1)} := F^{(m-1)} \amalg_{\Sigma[0,1] \amalg_{\Sigma[0,0]} \Sigma[1,0]} \Sigma[1,1],$$

which implies that the inclusion

$$F^{(m-1)} \hookrightarrow F^{(m-1,1)}$$

is an acyclic cofibration.

- (2) To obtain $F^{(m-1,2)}$ from $F^{(m-1,1)}$, we want to include all horizontal composites involving the new square. This step can be described as a pushout

$$F^{(m-1,2)} := F^{(m-1,1)} \amalg_{\Sigma[1,n-m-1] \amalg_{\Sigma[1,0]} \Sigma[1,1]} \Sigma[1,n-m],$$

and the inclusion

$$F^{(m-1,1)} \hookrightarrow F^{(m-1,2)}$$

is therefore an acyclic cofibration.

- (3) Similarly, we obtain $F^{(m-1,3)}$ from $F^{(m-1,2)}$ by adding all vertical composites involving the new square, a process given by the pushout

$$F^{(m-1,3)} := F^{(m-1,2)} \amalg_{\Sigma[m-1,1] \amalg_{\Sigma[0,1]} \Sigma[1,1]} \Sigma[m,1],$$

and the inclusion

$$F^{(m-1,2)} \hookrightarrow F^{(m-1,3)}$$

is again an acyclic cofibration.

- (4) Finally, we obtain $F^{(m)} := F^{(m-1,4)}$ from $F^{(m-1,3)}$ by gluing a rectangular grid to ensure compatibility of the two types of composition, as described in [Remark 2.4](#). This step can be described as a pushout of $F^{(m-1,3)}$ with $\Sigma[m, n - m]$ along their intersection. The inclusion

$$F^{(m-1,3)} \hookrightarrow F^{(m-1,4)}$$

is therefore an acyclic cofibration.

- Returning to our original filtration, we now obtain $F^{(n)}$ from $F^{(n-1)}$ by adding a new element to the augmentation and all vertical composites in the last column, as depicted in the codomain of the map j_4 in the diagram above when $n = 4$. This step is described by a pushout

$$F^{(n)} := F^{(n-1)} \amalg_{\Sigma[n-1,0]} V[n],$$

and therefore the inclusion of

$$F^{(n-1)} \hookrightarrow F^{(n)}$$

is an acyclic cofibration.

Finally, by direct verification one can check that $W[n] \cong F^{(n)}$, which concludes the proof. \square

Proof of [Lemma 5.9](#) We need to prove that the map $W[P] \rightarrow W[n]$ is an acyclic cofibration in $\mathcal{C}_J^{\Sigma^{\text{op}}}$. Since it is an inclusion, we need only show that it is a weak equivalence.

Consider the commutative diagram of inclusions

$$\begin{array}{ccc} H[P] & \hookrightarrow & W[P] \\ \downarrow & & \downarrow \\ H[n] & \hookrightarrow & W[n] \end{array}$$

in which the left-hand vertical map is a weak equivalence by [Lemma 5.10](#) and the bottom horizontal map is a weak equivalence by [Lemma 5.11](#). To prove our desired result, it suffices to prove that the top horizontal map is a weak equivalence.

We observed in [Definition 4.6](#) that $W[P]$ can be obtained as a pushout of $W[2]$ and $W[n-1]$ along $W[1]$, and similarly $H[P]$ can be obtained as a pushout of $H[2]$ and $H[n-1]$ along $H[1]$. Since these pushouts are taken along cofibrations, they are homotopy pushouts, and thus the weak equivalences of [Lemma 5.11](#) assemble into the upper horizontal map being a weak equivalence. \square

6 The Quillen equivalence

We are now ready to prove the main result of this paper, namely that the Quillen pair from [Theorem 5.1](#) is indeed a Quillen equivalence.

Theorem 6.1 *The path construction and the S_\bullet -construction induce a Quillen equivalence*

$$\mathcal{P}: \mathcal{C}_S^{\Delta\text{op}} \rightleftarrows \mathcal{C}_{\mathcal{T}}^{\Sigma\text{op}} : S_\bullet.$$

Our strategy for the proof is to show that

- the functor \mathcal{P} reflects weak equivalences, and
- for every fibrant object Y of $\mathcal{C}_{\mathcal{T}}^{\Sigma\text{op}}$, the counit of the map $\mathcal{P}S_\bullet Y \rightarrow Y$ is a weak equivalence.

These conditions constitute the characterization of Quillen equivalences from [\[21, Corollary 1.3.16\]](#) in the special case when all objects are cofibrant. The former statement appears in the next proposition, while the latter is given as [Proposition 6.5](#) below.

Proposition 6.2 *The path construction functor*

$$\mathcal{P}: \mathcal{C}_S^{\Delta\text{op}} \rightarrow \mathcal{C}_{\mathcal{T}}^{\Sigma\text{op}}$$

reflects weak equivalences.

Before embarking on the proof, we show that, while \mathcal{P} does not preserve fibrant objects, it does so up to injective fibrant replacement.

Notation 6.3 If Y is an object of $\mathcal{C}^{\Sigma\text{op}}$, we denote by Y^f a functorial fibrant replacement of Y in the injective model structure. In particular, the accompanying map $Y \rightarrow Y^f$ is a levelwise weak equivalence.

Lemma 6.4 *If X is a fibrant object of $\mathcal{C}_S^{\Delta\text{op}}$, then $(\mathcal{P}X)^f$ is fibrant in $\mathcal{C}_{\mathcal{T}}^{\Sigma\text{op}}$.*

Proof As $(\mathcal{P}X)^f$ is by definition injectively fibrant, we only need to prove that $(\mathcal{P}X)^f$ is \mathcal{T} -local. We verify that the map

$$\text{Map}^h(\Sigma[1, 1], (\mathcal{P}X)^f) \rightarrow \text{Map}^h(\Sigma[1, 0] \amalg_{\Sigma[0, 0]} \Sigma[0, 1], (\mathcal{P}X)^f),$$

induced by

$$\tau^{1,1}: \Sigma[1, 0] \amalg_{\Sigma[0, 0]} \Sigma[0, 1] \hookrightarrow \Sigma[1, 1],$$

as described in [Definition 4.4](#), is a weak equivalence. The other cases are similar; in particular, [Proposition 4.3](#) can be used to verify locality with respect to \mathcal{T}_{aug} .

Since X is \mathcal{S} -local, the 2-Segal map

$$X_3 \xrightarrow{(d_2, d_0)} X_2 \times_{X_1}^h X_2.$$

is a weak equivalence. Using the definition of the functor \mathcal{P} , we can rewrite this map as

$$(\mathcal{P}X)_{1,1} \rightarrow (\mathcal{P}X)_{1,0} \times_{(\mathcal{P}X)_{0,0}}^h (\mathcal{P}X)_{0,1}$$

induced by $\tau^{1,1}$. Since the injective fibrant replacement $(\mathcal{P}X)^f$ of $\mathcal{P}X$ is levelwise weakly equivalent to $\mathcal{P}X$, the map

$$(\mathcal{P}X)_{1,1}^f \rightarrow (\mathcal{P}X)_{1,0}^f \times_{(\mathcal{P}X)_{0,0}^f}^h (\mathcal{P}X)_{0,1}^f$$

is also a weak equivalence. But this weak equivalence can be identified with the map

$$\begin{array}{c} \text{Map}(\Sigma[1, 1], (\mathcal{P}X)^f) \\ \searrow \\ \text{Map}(\Sigma[1, 0], (\mathcal{P}X)^f) \times_{\text{Map}(\Sigma[0, 0], (\mathcal{P}X)^f)}^h \text{Map}(\Sigma[0, 1], (\mathcal{P}X)^f) \end{array}$$

which in turn is the map

$$\text{Map}(\Sigma[1, 1], (\mathcal{P}X)^f) \rightarrow \text{Map}(\Sigma[1, 0] \amalg_{\Sigma[0, 0]} \Sigma[0, 1], (\mathcal{P}X)^f)$$

induced by $\tau^{1,1}$. Because $(\mathcal{P}X)^f$ is injectively fibrant, this map models a weak equivalence on derived mapping spaces,

$$\text{Map}^h(\Sigma[1, 1], (\mathcal{P}X)^f) \rightarrow \text{Map}^h(\Sigma[1, 0] \amalg_{\Sigma[0, 0]} \Sigma[0, 1], (\mathcal{P}X)^f).$$

Hence $(\mathcal{P}X)^f$ is local with respect to this map, as desired. \square

Proof of Proposition 6.2 Suppose that $g: X \rightarrow X'$ is a map in $\mathcal{C}^{\Delta^{\text{op}}}$ such that $\mathcal{P}g: \mathcal{P}X \rightarrow \mathcal{P}X'$ is a weak equivalence in $\mathcal{C}_{\mathcal{T}}^{\Sigma^{\text{op}}}$. We want to show that g is a weak equivalence in $\mathcal{C}_{\mathcal{S}}^{\Delta^{\text{op}}}$.

Assume first that X and X' are \mathcal{S} -local. By Lemma 6.4, the injectively fibrant replacements $(\mathcal{P}X)^f$ and $(\mathcal{P}X')^f$ are \mathcal{T} -local. Moreover, $(\mathcal{P}g)^f$ is a weak equivalence in $\mathcal{C}_{\mathcal{T}}^{\Sigma^{\text{op}}}$. In particular, $(\mathcal{P}g)^f$ is a \mathcal{T} -local equivalence between \mathcal{T} -local objects, so it is a levelwise weak equivalence. In the commutative diagram

$$\begin{array}{ccc} \mathcal{P}X & \xrightarrow{\mathcal{P}g} & \mathcal{P}X' \\ \downarrow & & \downarrow \\ (\mathcal{P}X)^f & \xrightarrow{(\mathcal{P}g)^f} & (\mathcal{P}X')^f \end{array}$$

the vertical maps are also levelwise weak equivalences, so $\mathcal{P}g$ is must be as well.

Finally, given that $\mathcal{P}g$ is a levelwise weak equivalence, the map $g_0 = (\mathcal{P}g)_{-1}$ is a weak equivalence, and, similarly, for every n we get that $g_n = (\mathcal{P}g)_{n-1,0}$ is a weak equivalence. We conclude that g is a levelwise weak equivalence, and in particular an \mathcal{S} –local equivalence, as desired.

For arbitrary X and X' , let

$$(\widetilde{-})\colon \mathcal{C}_\mathcal{S}^{\Delta\mathrm{op}} \rightarrow \mathcal{C}_\mathcal{S}^{\Delta\mathrm{op}}$$

denote a functorial fibrant replacement. Then g and $\mathcal{P}g$ fit into commutative diagrams

$$\begin{array}{ccc} X & \xrightarrow{g} & X' \\ \simeq_\mathcal{S} \downarrow & & \downarrow \simeq_\mathcal{S} \\ \widetilde{X} & \xrightarrow{\widetilde{g}} & \widetilde{X}' \end{array} \rightsquigarrow \begin{array}{ccc} \mathcal{P}X & \xrightarrow{\mathcal{P}g} & \mathcal{P}X' \\ \simeq_\mathcal{T} \downarrow & & \downarrow \simeq_\mathcal{T} \\ \mathcal{P}\widetilde{X} & \xrightarrow{\mathcal{P}\widetilde{g}} & \mathcal{P}\widetilde{X}' \end{array}$$

where we use from [Theorem 5.1](#) that the image of a weak equivalence in $\mathcal{C}_\mathcal{T}^{\Sigma\mathrm{op}}$ under \mathcal{P} is a weak equivalence in $\mathcal{C}_\mathcal{S}^{\Delta\mathrm{op}}$.

Suppose that $\mathcal{P}g$ is a weak equivalence in $\mathcal{C}_\mathcal{T}^{\Sigma\mathrm{op}}$. Then $\mathcal{P}\widetilde{g}$ is also a weak equivalence in $\mathcal{C}_\mathcal{T}^{\Sigma\mathrm{op}}$. By the previous argument, \widetilde{g} is a weak equivalence in $\mathcal{C}_\mathcal{S}^{\Delta\mathrm{op}}$, from which we can conclude that g is also, as desired. □

It remains to prove the aforementioned condition on the counit.

Proposition 6.5 *If Y is a fibrant object of $\mathcal{C}_\mathcal{T}^{\Sigma\mathrm{op}}$, then the counit map*

$$\epsilon_Y\colon \mathcal{P}S_\bullet Y \rightarrow Y$$

is a levelwise weak equivalence, and therefore a weak equivalence in $\mathcal{C}_\mathcal{T}^{\Sigma\mathrm{op}}$.

We need a preliminary result, which roughly says that, in the \mathcal{T} –localized model structure, $\mathsf{W}[q+1+r]$ is weakly equivalent to the copy of $\Sigma[q,r]$ contained inside of it. The argument is similar to that of [Lemma 5.11](#).

Lemma 6.6 *The inclusion $\Sigma[q,r] \hookrightarrow \mathsf{W}[q+1+r]$ is an acyclic cofibration in $\mathcal{C}_\mathcal{T}^{\Sigma\mathrm{op}}$.*

Proof This inclusion fits into a commutative diagram of inclusions of the form

$$\begin{array}{ccc} \Sigma[q,r] & \hookrightarrow & \mathsf{W}[q+1+r] \\ \uparrow & & \uparrow \\ \Sigma[q,0] \amalg_{\Sigma[0,0]} \Sigma[0,r] & \hookrightarrow & V[q+1] \amalg_{\Sigma[0,0]} \Sigma[0,r] \hookrightarrow \mathsf{W}[q+1] \amalg_{\Sigma[0,0]} \Sigma[0,r] \end{array}$$

The left-hand vertical map is an element of $\mathcal{T}_{\text{stable}}$. The left-most bottom horizontal map is a pushout along the map $\Sigma[q, 0] \hookrightarrow V[q + 1]$ in \mathcal{T}_{aug} . The right-most bottom horizontal map is a pushout along the map $V[q + 1] \hookrightarrow W[q + 1]$, which can be proven to be an acyclic cofibration by dualizing the argument from [Lemma 5.11](#). The right-hand vertical map fits into a commutative diagram of inclusions of the form

$$\begin{array}{ccc} H[q + 1] \amalg_{\Sigma[0, 0]} \Sigma[0, r] & \hookrightarrow & H[q + 1 + r] \\ \downarrow & & \downarrow \\ W[q + 1] \amalg_{\Sigma[0, 0]} \Sigma[0, r] & \hookrightarrow & W[q + 1 + r] \end{array}$$

where the two vertical maps are weak equivalences by [Lemma 5.11](#) and the top map can be written as a pushout along a map in $\mathcal{T}_{\text{Segal}}$. \square

Proof of Proposition 6.5 We first observe that the precomposition functor $\mathcal{P} = p^*$ admits a \mathcal{C} –enriched left adjoint

$$p_! : \mathcal{C}^{\Sigma^{\text{op}}} \rightarrow \mathcal{C}^{\Delta^{\text{op}}},$$

which is given by left Kan extension [[25](#), Theorem 4.50]. Since left Kan extensions are well behaved on representables [[30](#), Section 7.7], there are isomorphisms

$$p_! \Sigma[q, r] \cong \Delta[q + 1 + r] \quad \text{and} \quad p_! \Sigma[-1] \cong \Delta[0].$$

The counit map evaluated at (q, r) ,

$$\epsilon_Y : (p^* p_* Y)_{q, r} \rightarrow Y_{q, r},$$

can be identified with a map

$$\text{Map}_{\mathcal{C}^{\Sigma^{\text{op}}}}(\Sigma[q, r], p^* p_* Y) \rightarrow \text{Map}_{\mathcal{C}^{\Sigma^{\text{op}}}}(\Sigma[q, r], Y)$$

via the enriched Yoneda lemma. Using the fact that $\mathcal{P} = p^*$ and $S_\bullet = p_*$ are right adjoints (of $p_!$ and p^* , respectively) the left-hand side in the above map can be identified with

$$\text{Map}_{\mathcal{C}^{\Sigma^{\text{op}}}}(\Sigma[q, r], p^* p_* Y) \cong \text{Map}_{\mathcal{C}^{\Sigma^{\text{op}}}}(p^* p_! \Sigma[q, r], Y).$$

Applying [Remark 2.17](#) and our adjunctions, we have that

$$W[q + 1 + r] \cong \mathcal{P} \Delta[q + 1 + r] \cong \mathcal{P} p_! \Sigma[q, r] = p^* p_! \Sigma[q, r],$$

hence the mapping space can further be identified with

$$\text{Map}_{\mathcal{C}^{\Sigma^{\text{op}}}}(p^* p_! \Sigma[q, r], Y) \cong \text{Map}_{\mathcal{C}^{\Sigma^{\text{op}}}}(W[q + 1 + r], Y)$$

and the map above can be identified with

$$\mathrm{Map}_{\mathcal{C}^{\Sigma^{\mathrm{op}}}}(\mathbf{W}[q+1+r], Y) \rightarrow \mathrm{Map}_{\mathcal{C}^{\Sigma^{\mathrm{op}}}}(\Sigma[q, r], Y)$$

induced by the inclusion $\Sigma[q, r] \hookrightarrow \mathbf{W}[q+1+r]$. Since Y is fibrant in $\mathcal{C}_J^{\Sigma^{\mathrm{op}}}$, this map models the analogous map on derived mapping spaces, which is a weak equivalence by [Lemma 6.6](#). The -1 -component can be treated similarly. \square

7 Variants of the model structures

We conclude with some illustrations that the techniques employed in the paper are quite flexible and can be used to obtain further results along the same lines. Indeed, by performing enriched localizations with respect to different sets of maps, or by localizing the projective model structure instead of the injective model structure, we can obtain variants of [Theorem 6.1](#).

7.1 Alternative localizations

In applications one might have further properties on 2-Segal objects and might wonder how to translate them to the context of augmented double Segal objects. We consider two natural such conditions here.

Reduced and pointed settings While the augmentation condition for stable double Segal objects might seem complicated, it is intended to serve as an appropriate generalization of zero objects. For augmented stable double Segal objects coming from *pointed* contexts, such as those from [Section 3.2](#), the augmentation represents a space of zero objects, and is therefore contractible. We can restrict ourselves to this setting as follows.

Definition 7.1 A *stable pointed double Segal object* in \mathcal{C} is an augmented stable double Segal object Y such that the map

$$Y_{-1} \rightarrow *$$

to the terminal object is a weak equivalence in \mathcal{C} .

For 2-Segal objects, the analogous property was considered in [\[6, Definition 1.9\]](#) in the case where $\mathcal{C} = \mathbf{Set}$, but it can be generalized as follows.

Definition 7.2 A simplicial object X in \mathcal{C} is *reduced* if the map from X_0 to the terminal object in \mathcal{C} is a weak equivalence in \mathcal{C} .

By means of [Theorem 1.11](#), there are model structures on $\mathcal{C}^{\Delta^{\text{op}}}$ and on $\mathcal{C}^{\Sigma^{\text{op}}}$ whose fibrant objects are reduced 2–Segal objects and pointed stable double Segal objects, respectively.

Segal objects We know that every Segal object is 2–Segal, so we can ask how to identify the corresponding preaugmented bisimplicial objects. In the context of reduced Segal objects, as just discussed, they can be described in terms of homological algebra, as we now explain.

Let us consider the following condition. Given an augmented stable double Segal object Y , there is a natural notion of an *object of extensions* in Y , in the form of distinguished squares in which the left bottom corner is a zero object, realized as a homotopy pullback

$$Y_{1,1} \overset{h}{\times}_{Y_{0,0}} Y_{-1}.$$

This definition recovers, in particular, the groupoid of exact sequences in an exact category, as well as the space of fiber sequences in a stable $(\infty, 1)$ –category.

As in homological algebra, a natural question is whether all the extensions in Y split. We thus make the following definition:

Definition 7.3 A stable pointed double Segal object Y in \mathcal{C} is *split* if the map

$$(s_v \circ s_h \circ \text{pr}_1, t_v \circ t_h \circ \text{pr}_1): Y_{1,1} \overset{h}{\times}_{Y_{0,0}} Y_{-1} \rightarrow Y_{0,0} \times Y_{0,0}$$

is a weak equivalence.

Remark 7.4 Let us give some heuristic motivation for this definition. We would like to encode the structure of $Y = \mathcal{P}(X)$ for a given Segal object X . It is not hard to show that a 2–Segal object is Segal if and only if the Segal map $X_2 \xrightarrow{(d_2, d_0)} X_1 \overset{h}{\times}_{X_0} X_1$ is an equivalence. While we would like to translate this condition to Y , the maps d_2 and d_0 have different domains there (namely, $d_2: Y_{0,1} \rightarrow Y_{0,0}$ and $d_0: Y_{1,0} \rightarrow Y_{0,0}$) and hence cannot be combined to a single map. Instead, we use the following reformulation: since X is reduced, we have a weak equivalence

$$X_3 \overset{h}{\times}_{X_1} X_0 \xrightarrow{\sim} X_2.$$

Hence, we can identify the map from [Definition 7.3](#) in this case with

$$X_3 \overset{h}{\times}_{X_1} X_0 \rightarrow X_1 \overset{h}{\times}_{X_0} X_1$$

and require it to be a weak equivalence.

Proposition 7.5 The generalized S_\bullet –construction induces Quillen equivalences

$$\mathcal{P}: \mathcal{C}^{\Delta^{\text{op}}} \rightleftarrows \mathcal{C}^{\Sigma^{\text{op}}} : S_\bullet.$$

- between the model structure for reduced 2–Segal objects and the model structure for stable pointed double Segal objects; and
- between the model structure for reduced Segal objects and the model structure for split stable pointed double Segal objects.

7.2 Localizations of the projective model structure

The examples of augmented stable double Segal spaces provided in Sections 3.1 and 3.2 are levelwise Kan complexes, and in particular fibrant in the projective model structure. Since it is a stronger condition to be injectively fibrant, working in that context presents a potential obstruction to finding natural examples of \mathcal{T} –local objects. As an example coming from bordisms, one can construct a homotopical version of the 2–Segal set of bordisms with genus constraints from [6, Examples 2.2 and 7.2] using methods from [9].

For this reason, we also want a model structure for augmented stable double Segal objects obtained as a localization of the projective model structure, so that the fibrant objects are precisely the levelwise fibrant augmented stable double Segal objects. We can obtain the same kind of variant to get levelwise fibrant 2–Segal objects.

By an analogue of [20, Theorem 3.3.20], the identity functors realize Quillen equivalences

$$\mathrm{id} \colon \mathcal{C}_{\mathrm{proj},S}^{\Delta^{\mathrm{op}}} \rightleftarrows \mathcal{C}_{\mathrm{inj},S}^{\Delta^{\mathrm{op}}} \colon \mathrm{id} \quad \text{and} \quad \mathrm{id} \colon \mathcal{C}_{\mathrm{proj},S}^{\Sigma^{\mathrm{op}}} \rightleftarrows \mathcal{C}_{\mathrm{inj},S}^{\Sigma^{\mathrm{op}}} \colon \mathrm{id}.$$

Combining these equivalences with the ones from our main result, we obtain the following comparisons:

Proposition 7.6 *There is a zigzag of Quillen equivalences*

$$\begin{array}{ccc} \mathcal{C}_{\mathrm{inj},S}^{\Delta^{\mathrm{op}}} & \begin{array}{c} \xleftarrow{\mathcal{P}} \\ \xrightarrow{S_{\bullet}} \end{array} & \mathcal{C}_{\mathrm{inj},\mathcal{T}}^{\Sigma^{\mathrm{op}}} \\ \mathrm{id} \updownarrow \mathrm{id} & & \mathrm{id} \updownarrow \mathrm{id} \\ \mathcal{C}_{\mathrm{proj},S}^{\Delta^{\mathrm{op}}} & & \mathcal{C}_{\mathrm{proj},\mathcal{T}}^{\Sigma^{\mathrm{op}}} \end{array}$$

References

[1] **D W Anderson**, *Fibrations and geometric realizations*, Bull. Amer. Math. Soc. 84 (1978) 765–788 [MR](#) [Zbl](#)

[2] **C Barwick**, (∞, n) –Cat as a closed model category, PhD thesis, University of Pennsylvania (2005) Available at <https://search.proquest.com/docview/305445747>

- [3] **C Barwick**, *On left and right model categories and left and right Bousfield localizations*, Homology Homotopy Appl. 12 (2010) 245–320 [MR](#) [Zbl](#)
- [4] **C Barwick**, *On the Q construction for exact ∞ –categories*, preprint (2013) [arXiv](#)
- [5] **C Barwick**, *On the algebraic K –theory of higher categories*, J. Topol. 9 (2016) 245–347 [MR](#) [Zbl](#)
- [6] **J E Bergner, A M Osorno, V Ozornova, M Rovelli, C I Scheimbauer**, *2–Segal sets and the Waldhausen construction*, Topology Appl. 235 (2018) 445–484 [MR](#) [Zbl](#)
- [7] **J E Bergner, A M Osorno, V Ozornova, M Rovelli, C I Scheimbauer**, *Comparison of Waldhausen constructions*, preprint (2019) [arXiv](#)
- [8] **A J Blumberg, D Gepner, G Tabuada**, *A universal characterization of higher algebraic K –theory*, Geom. Topol. 17 (2013) 733–838 [MR](#) [Zbl](#)
- [9] **D Calaque, C Scheimbauer**, *A note on the (∞, n) –category of cobordisms*, Algebr. Geom. Topol. 19 (2019) 533–655 [MR](#) [Zbl](#)
- [10] **L Carlier**, *Incidence bicomodules, Möbius inversion and a Rota formula for infinity adjunctions*, Algebr. Geom. Topol. 20 (2020) 169–213 [MR](#) [Zbl](#)
- [11] **T Dyckerhoff, M Kapranov**, *Higher Segal spaces*, Lecture Notes in Math. 2244, Springer (2019) [MR](#) [Zbl](#)
- [12] **M Feller, R Garner, J Kock, M U Proulx, M Weber**, *Every 2–Segal space is unital*, Commun. Contemp. Math. 23 (2021) art. id. 2050055 [MR](#) [Zbl](#)
- [13] **T M Fiore, S Paoli**, *A Thomason model structure on the category of small n –fold categories*, Algebr. Geom. Topol. 10 (2010) 1933–2008 [MR](#) [Zbl](#)
- [14] **T M Fiore, S Paoli, D Pronk**, *Model structures on the category of small double categories*, Algebr. Geom. Topol. 8 (2008) 1855–1959 [MR](#) [Zbl](#)
- [15] **T M Fiore, M Pieper**, *Waldhausen additivity: classical and quasicategorical*, J. Homotopy Relat. Struct. 14 (2019) 109–197 [MR](#) [Zbl](#)
- [16] **I Gálvez-Carrillo, J Kock, A Tonks**, *Decomposition spaces, incidence algebras and Möbius inversion, I: Basic theory*, Adv. Math. 331 (2018) 952–1015 [MR](#) [Zbl](#)
- [17] **M Grandis, R Pare**, *Limits in double categories*, Cahiers Topologie Géom. Différentielle Catég. 40 (1999) 162–220 [MR](#) [Zbl](#)
- [18] **B J Guillou, J P May, M Merling, A M Osorno**, *Symmetric monoidal G –categories and their strictification*, 1 (2020) [MR](#) [Zbl](#)
- [19] **R Haugseng**, *Iterated spans and classical topological field theories*, Math. Z. 289 (2018) 1427–1488 [MR](#) [Zbl](#)
- [20] **P S Hirschhorn**, *Model categories and their localizations*, Math. Surv. Monogr. 99, Amer. Math. Soc., Providence, RI (2003) [MR](#) [Zbl](#)
- [21] **M Hovey**, *Model categories*, Math. Surv. Monogr. 63, Amer. Math. Soc., Providence, RI (1999) [MR](#) [Zbl](#)
- [22] **L Illusie**, *Complexe cotangent et déformations, II*, Lecture Notes in Math. 283, Springer (1972) [MR](#) [Zbl](#)

- [23] **A Joyal**, *Notes on quasi-categories*, preprint (2008) Available at <https://tinyurl.com/joyalpre>
- [24] **A Joyal, M Tierney**, *Quasi-categories vs Segal spaces*, from “Categories in algebra, geometry and mathematical physics” (A Davydov, M Batanin, M Johnson, S Lack, A Neeman, editors), Contemp. Math. 431, Amer. Math. Soc., Providence, RI (2007) 277–326 [MR](#) [Zbl](#)
- [25] **G M Kelly**, *Basic concepts of enriched category theory*, Lond. Math. Soc. Lect. Note Ser. 64, Cambridge Univ. Press (1982) [MR](#) [Zbl](#)
- [26] **J Lurie**, *Stable infinity categories*, preprint (2006) [arXiv](#)
- [27] **J Lurie**, *Higher topos theory*, Ann. of Math. Stud. 170, Princeton Univ. Press (2009) [MR](#) [Zbl](#)
- [28] **D G Quillen**, *Homotopical algebra*, Lecture Notes in Math. 43, Springer (1967) [MR](#) [Zbl](#)
- [29] **D Quillen**, *Higher algebraic K-theory, I*, from “Algebraic K-theory, I: Higher K-theories” (H Bass, editor), Lecture Notes in Math. 341, Springer (1973) 85–147 [MR](#) [Zbl](#)
- [30] **E Riehl**, *Categorical homotopy theory*, New Math. Monogr. 24, Cambridge Univ. Press (2014) [MR](#) [Zbl](#)
- [31] **B Toën, G Vezzosi**, *Homotopical algebraic geometry, I: Topos theory*, Adv. Math. 193 (2005) 257–372 [MR](#) [Zbl](#)
- [32] **F Waldhausen**, *Algebraic K-theory of spaces*, from “Algebraic and geometric topology” (A Ranicki, N Levitt, F Quinn, editors), Lecture Notes in Math. 1126, Springer (1985) 318–419 [MR](#) [Zbl](#)

JEB: Department of Mathematics, University of Virginia
Charlottesville, VA, United States

AMO: Department of Mathematics, Reed College
Portland, OR, United States

VO: Fakultät für Mathematik, Ruhr-Universität Bochum
Bochum, Germany

MR: Mathematical Sciences Institute, The Australian National University
Canberra, ACT, Australia

Current address: Department of Mathematics and Statistics, University of Massachusetts
Amherst, MA, United States

CIS: Zentrum Mathematik, Technische Universität München
Garching bei München, Germany

jeb2md@virginia.edu, aosorno@reed.edu, viktoriya.ozornova@rub.de,
rovelli@math.umass.edu, scheimbauer@tum.de

Received: 25 January 2019 Revised: 3 February 2020

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

MANAGING EDITOR

Colin Rourke
agt@msp.warwick.ac.uk
University of Warwick

BOARD OF EDITORS

Matthew Ando	University of Illinois mando@math.uiuc.edu	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Joan Birman	Columbia University jb@math.columbia.edu	Dan Margalit	Georgia Institute of Technology margalit@math.gatech.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Norihiko Minami	Nagoya Institute of Technology nori@nitech.ac.jp
Fred Cohen	University of Rochester cohf@math.rochester.edu	Andrés Navas Flores	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Thomas Nikolaus	University of Münster nikolaus@uni-muenster.de
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Robert Oliver	Université Paris-Nord bobol@math.univ-paris13.fr
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Luis Paris	Université de Bourgogne lparis@u-bourgogne.fr
David Futer	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
Soren Galatius	Stanford University galatius@math.stanford.edu	Peter Scott	University of Michigan pscott@umich.edu
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Zoltán Szabó	Princeton University szabo@math.princeton.edu
J. Elisenda Grigsby	Boston College grigsbyj@bc.edu	Ulrike Tillmann	Oxford University tillmann@maths.ox.ac.uk
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Daniel T. Wise	McGill University, Canada daniel.wise@mcgill.ca
Tsuyoshi Kobayashi	Nara Women's University tsuyoshi09@gmail.com		


See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2021 is US\$560/year for the electronic version, and \$835/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 7 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 21 Issue 3 (pages 1075–1593) 2021

On the Upsilon invariant of cable knots	1075
WENZHAO CHEN	
A–infinity algebras, strand algebras, and contact categories	1093
DANIEL V MATHEWS	
Barcode embeddings for metric graphs	1209
STEVE OUDOT and ELCHANAN SOLOMON	
2–Segal objects and the Waldhausen construction	1267
JULIA E BERGNER, ANGÉLICA M OSORNO, VIKTORIYA OZORNOVA, MARTINA ROVELLI and CLAUDIA I SCHEIMBAUER	
Contractible open manifolds which embed in no compact, locally connected and locally 1–connected metric space	1327
SHIJIE GU	
Limits of sequences of pseudo-Anosov maps and of hyperbolic 3–manifolds	1351
SYLVAIN BONNOT, ANDRÉ DE CARVALHO, JUAN GONZÁLEZ-MENESES and TOBY HALL	
Homological stability for moduli spaces of disconnected submanifolds, I	1371
MARTIN PALMER	
Configuration spaces of squares in a rectangle	1445
LEONID PLACHTA	
Coloring invariants of knots and links are often intractable	1479
GREG KUPERBERG and ERIC SAMPERTON	
Powers of Dehn twists generating right-angled Artin groups	1511
DONGGYUN SEO	
Rational homotopy equivalences and singular chains	1535
MANUEL RIVERA, FELIX WIERSTRA and MAHMOUD ZEINALIAN	
A combinatorial description of the centralizer algebras connected to the Links–Gould invariant	1553
CRISTINA ANA-MARIA ANGHEL	